



**Report of Rapid Biodiversity Assessments at
Shiwandashan National Nature Reserve and
National Forest Park, Southwest Guangxi, China,
2000 and 2001**

Kadoorie Farm and Botanic Garden
in collaboration with
Guangxi Forestry Department
Guangxi Institute of Botany
South China Normal University

July 2003

South China Forest Biodiversity Survey Report Series: No. 35
(Online Simplified Version)

Report of Rapid Biodiversity Assessments at Shiwandashan National Nature Reserve and National Forest Park, Southwest Guangxi, China, 2000 and 2001

Editors

John R. Fellowes, Bosco P.L. Chan, Michael W.N. Lau, Ng Sai-Chit and Gloria L.P. Siu

Contributors

Kadoorie Farm and Botanic Garden:	Billy C.H. Hau	(BH)
	Lee Kwok Shing	(LKS)
	Michael W.N. Lau	(ML)
	John R. Fellowes	(JRF)
	Ng Sai-Chit	(NSC)
	Gloria L.P. Siu	(GS)
Bosco P.L. Chan		(BC)
Guangxi Forestry Department:	Xu Zhihong	(XZH)
	Xu Gesheng	(XGS)
Guangxi Institute of Botany:	Li Guangzhao	(LGZ)
	Tang Saichun	(TSC)
Institute of Botany, Chinese Academy of Sciences	Jin Xiaohua	(JXH)
South China Normal University:	Xiao Zhi	(XZ)
	Chen Xianglin	(CXL)
Chinese Academy of Forestry:	Jiang Jingman	(JJM)
	Li Yide	(LYD)
	Liu Fanglin	(LFL)
Guangxi University:	Zhou Fang	(ZF)
Guangxi Normal University:	Zhou Shanyi	(ZSY)
Xinyang Teachers' College:	Li Hongjing	(LHJ)
Station Biologique, La Tour du Valat:	Heinz Hafner	(HH)
Voluntary specialists:	Graham T. Reels	(GTR)
	Keith D.P. Wilson	(KW)

Background

The present report details the findings of visits to Southwest Guangxi by members of Kadoorie Farm and Botanic Garden (KFBG) in Hong Kong and their colleagues, as part of KFBG's South China Biodiversity Conservation Programme. The overall aim of the programme is to minimise the loss of forest biodiversity in the region, and the emphasis in the first phase is on gathering up-to-date information on the distribution and status of fauna and flora.

Citation

Kadoorie Farm and Botanic Garden, 2003. *Report of Rapid Biodiversity Assessments at Shiwandashan National Nature Reserve and National Forest Park, Southwest Guangxi, China, 2000 and 2001*. South China Forest Biodiversity Survey Report Series (Online Simplified Version): No. 35. KFBG, Hong Kong SAR, ii + 30 pp.

Copyright

© Kadoorie Farm and Botanic Garden Corporation
Lam Kam Road, Tai Po, N.T., Hong Kong SAR

July 2003

Contents

Objectives	1
Methods	1
Location and management	2
Results	2
<i>Vegetation</i>	2
<i>Flora</i>	3
<i>Mammals</i>	14
<i>Birds</i>	19
<i>Reptiles and Amphibians</i>	21
<i>Fish</i>	22
<i>Ants</i>	23
<i>Dragonflies</i>	24
<i>Butterflies</i>	24
Summary of flora and fauna	25
Threats and problems	26
Opportunities	26
Acknowledgements	27
References	27
Figure 1 Map.....	30

Translation of common Chinese geographical terms

Romanized Chinese (pinyin)	English meaning
Bei	north
Dao	island
Dong	east
Feng shui	the Chinese system of geomancy
Feng, Ding	peak
Gang	harbour
Hai	sea
He, Chuan, Jiang	river
Hu, Chi	lake
Keng, Gu, Gou	valley, stream
Kou	outlet
Ling	range
Nan	south
Ping	flat
Shan	mountain
Shi	city
Tun	hamlet
Wan	bay
Xi	west
Xi, Yong	stream
Xian	county
Xiang, Cun	village

Report of Rapid Biodiversity Assessments at Shiwandashan National Nature Reserve and National Forest Park, Southwest Guangxi, China, 2000 and 2001

Objectives

- The first trip was part of a species-specific survey for White-eared Night Heron *Gorsachius magnificus*, following past reports of the species at Shiwandashan (Zhou, 1996). The aims of the second survey were to collect up-to-date information on the fauna and flora of Shiwandashan National Nature Reserve and National Forest Park, and to use this to help determine conservation priorities within South China.

Methods

- On 31 March 2000 a team of biologists from KFBG (JRF, ML, LKS, Vicky Lam), France (HH), Guangxi (ZF, XZH, XGS) and Guangdong (Wang Ruijiang of South China Institute of Botany) visited Shiwandashan National Nature Reserve as part of a survey for White-eared Night Heron *Gorsachius magnificus* (Fellowes *et al.*, 2001).
- On 24 September 2000 a team from KFBG (ML, BH, GS, LKS, NSC, Winky Huen), Beijing (JJM, LYD, LFL), Guangxi (XZH, WFN, TSC, ZSY), Guangdong (XZ) and Henan (LHJ) visited Shiwandashan Nature Reserve. They arrived at Shangsi County Town at 15.45 after a rapid survey in Damingshan National Nature Reserve (Kadoorie Farm and Botanic Garden, 2003).
- On 28 September the team departed Shiwandashan National Forest Park for Nanning.
- On 7 December 2001 a team from KFBG (LKS, NSC), and Beijing (JXH) visited Shiwandashan National Nature Reserve at the invitation of Guangxi Forestry Department. They arrived at Shangsi County Town at 19.00 and met XZH.
- On 15 December at 09.30 they departed Shiwandashan National Forest Park and at 12.15 they were back at Nanning.
- During fieldwork visual searching for plants, mammals, birds, reptiles, amphibians, fish, ants, butterflies and dragonflies were conducted. Frogs and birds were also identified by their calls. Plant records were made by field observation, with some specimens collected.
- Status of large and medium-sized mammals (excluding Insectivora, Chiroptera and Muridae) at Shiwandashan was inferred largely based on interviews with local people in 1997 (Fellowes & Hau, 1997), with reference to colour pictures. For purposes of these interviews a list of South China mammals was compiled from various sources including Guangdong Forestry Department and South China Institute of Endangered Animals (1987), Corbet and Hill (1992) and Zhang Y. *et al.* (1997). The result is supplemented by additional data of Guangxi Forestry Survey and Planning Institute and Shiwandashan Nature Reserve (2002).
- Vascular plant records were made by LGZ and NSC and edited by NSC, except for orchids, for which records were made by JXH and GS and edited by GS. Mammal records were made by LKS, ML, JRF or BH. Records of birds were made or verified by LKS, ZF, HH or ML, reptiles and amphibians by ML, fish by BC and CXL, ants by JRF or ZS, dragonflies by KW, ML or GTR and butterflies by ML or GTR.
- Nomenclature in the report is standardised based, unless otherwise stated, on the following references:
 - Flora (Pteridophyta, Gymnospermae and Angiospermae excluding Orchidaceae): Anon. (1959-2001); Anon. (1991); Anon. (1996-2001); Anon. (2002a, 2002b); The Plant Names Project (2002);
 - Orchids (Angiospermae: Orchidaceae): Chen (1999); Lang (1999); Tsi (1999); Vogel & Turner (1992);
 - Mammals (Mammalia): D.E. Wilson & Cole (2000);
 - Birds (Aves): Inskipp *et al.* (1996);
 - Reptiles and Amphibians (Reptilia and Amphibia): Zhao E.-M. and Adler (1993); Zhao E. *et al.* (2000);

- Fish (Actinopterygii): Nelson (1994); Wu *et al.* (1999);
- Ants (Insecta: Hymenoptera: Formicidae): named species according to Bolton (1995); unnamed species with reference numbers according to the collection currently held by KFBG.
- Dragonflies (Insecta: Odonata): Schorr *et al.* (2001a, 2001b);
- Butterflies (Insecta: Lepidoptera): Bascombe (1995).
- Information on the global status of species is from IUCN publications, notably IUCN (2002). Certain taxa, including orchids, reptiles, amphibians, fish and invertebrates, have yet to be properly assessed for global status. National conservation status of orchids is based on Wang *et al.* (in press).
- Protected status in China is based on Hua and Yan (1993) for animals, and State Forestry Administration & Ministry of Agriculture (1999) for plants.

Location and management

- Shiwandashan National Nature Reserve is at the intersection of Shangsi, Fangcheng and Qinzhou Counties, southwest Guangxi, at 21°40'03"-22°04'18"N by 107°29'59"-108°13'11"E. The size of the nature reserve has been cited as 1,745 km² (MacKinnon *et al.*, 1996; Zhang W., 1998) but a figure of 583 km² is given by Guangxi Forestry Survey and Planning Institute (2002).
- The reserve has a relatively gentle mountainous landscape, with 82 peaks over 1,000 m. Altitude in the reserve ranges from below 300 m to 1,462 m at the summit of Shuliangling (also known as Jigongling). The geology is dominated by sandy shale, shale, conglomerate and granite (Guangxi Forestry Survey and Planning Institute and Shiwandashan Nature Reserve Management Station, 2002).
- The area has a tropical monsoon climate. The mean annual temperature is around 21°C. Mean monthly temperature ranges from around 13°C in January to 28°C in July. Annual precipitation in the reserve reaches 2,700 mm (Guangxi Forestry Survey and Planning Institute, 2002). Streams on the northern slopes drain to the Ming Jiang, a tributary of the Xi Jiang in the Zhujiang drainage system; streams on the southern slopes flow directly into the Gulf of Tonkin, which is also known as Beibu Wan (Zhao and Zhang, 2001).
- Shiwandashan National Nature Reserve established a reserve station in Shangsi County with 10 staff; two forestry police stations with 13 forestry police; and seven management stations with 80 staff. The area is densely populated with 11 villages and over 220,000 residents, of whom three-quarters live in Fangcheng County (Guangxi Forestry Survey and Planning Institute & Shiwandashan Nature Reserve Management Station, 2002).
- Shiwandashan was designated a provincial nature reserve in 1982 to protect its water catchment forest (MacKinnon *et al.*, 1996). It is classified as a Forest Ecosystem nature reserve (Zhang W., 1998). The nature reserve was upgraded to National-level in 2002 (Su Y., Guangxi Forestry Department, pers. comm., March 2003).

Results

Vegetation

- The zonal vegetation of the Shiwandashan region should be northern tropical monsoon rainforest. Major vegetation types in the region included monsoon rainforest, ravine rainforest, montane evergreen forest, montane dwarf forest, young secondary pine forest, and shrubland and grassland. Plantations of *Illicium verum* (Star Anise), *Pinus* spp., and *Cunninghamia lanceolata* (China fir) are also important landscape features of Shiwandashan (Forestry Department of Guangxi Zhuang Autonomous Region, 1993; Guangxi Forestry Survey and Planning Institute & Shiwandashan Nature Reserve, 2002).
- The present surveys visited only a limited area of the whole reserve. The vegetation of the surveyed area are mainly fragmented secondary forest in a matrix of degraded hillside shrubland, grassland, plantation and lowland agricultural land. The present surveys found extensive but fragmented cover of evergreen forest at Shiwandashan National Forest Park, Pinglongshan, Taiping and Sanliao Sidui.
- At Sanliao Sidui, the survey team visited secondary forest up to 15-25 m tall, and with trees up to 40 cm dbh. Such forest are very diverse and co-dominated largely by *Castanopsis* spp., *Lithocarpus* spp.,

Engelhardtia roxburghiana, *Eberhardtia aurata*, *Madhuca pasquieri*, *Pygeum topengii*, *Sarcosperma laurinum*, *Helicia longipetiolata*, *Elaeocarpus* spp. and *Artocarpus styracifolius*.

- At Taiping, lowland areas are mainly young secondary or remnant forests about 10-30 m tall, with trees less than 50 cm dbh, and dominated largely by *Castanopsis* spp., *Engelhardtia roxburghiana*, *Schima bambusifolia*, *Liquidambar formosana*, *Garcinia multiflora*, *Cylindrokelupha kerrii* and *Cryptocarya chingii*. Montane forest up to 10-15 m tall, with trees less than 50 cm dbh, was found at higher altitude. The forest was dominated by *Exbucklandia tonkinensis*, *Castanopsis carlesii*, *Lithocarpus* spp., *Rhododendron haofui*, *Hartia villosa*, *Diplopanax stachyanthus* and *Dendropanax hainanensis*.
- At Shiwandashan National Forest Park, secondary forest about 15-20 m tall, with trees up to 40 cm dbh, was found. The lowland forest was largely dominated by *Engelhardtia roxburghiana*, *Cryptocarya concinna*, *Cylindrokelupha tonkinensis*, *Pithecellobium clypearia*, *Macaranga sampsoni* and *M. henryi*. At higher altitude, the hillside forest was dominated by *Engelhardtia roxburghiana*, *Castanopsis fissa*, *C. hystrix*, *Mytilaria laosensis*, *Acronychia pedunculata*, *Litsea lancilimba* and *Michelia foveolata*.
- At Pinglongshan, secondary tropical lowland forest up to 10-30 m tall and with trees up to 50 cm dbh was found. The forest is largely dominated by *Castanopsis carlesii*, *Schima superba*, *Lithocarpus fenestrata*, *Helicia longipetiolata*, *Eberhardtia aurata*, *Cinnamomum porrectum*, and *Meliosma rigida*, and *Hopea chinensis*. The understorey is dominated by *Urophyllum chinensis*, *Pinanga discolor* and *Caryota monostachys*.

Flora

- The present surveys recorded 517 vascular plant species, including 59 fern species in 29 families, six gymnosperms in four families, and 452 flowering plant species in 95 families (Tables 1 and 2). This is a moderately high figure in altogether 9 days of fieldwork suggesting that the area had a diverse flora. The recorded flora is especially rich in tropical families and genera and hence is rather distinct from other forest vegetation in Guangxi.
- A number of new records of Guangxi were also found in the present surveys, including:
 - *Mecodium excertum*; a few colonies of this species were found at Shiwandashan National Forest Park.
 - *Euonymus mitratus* was locally common at Taiping and a few plants were also seen at Shiwandashan National Forest Park. It was previously recorded from South Yunnan, Vietnam and Cambodia.
 - *Vaccinium chunii* has only been recorded in Hainan, but a single plant was noticed.
 - *Carex tenuispicula* was previously known from Fujian and Guangdong. A small colony was found.
 - *Microgonium beccarianum* is a new record for China mainland and had only been recorded in Taiwan and some Pacific islands. A small colony was noticed but it is likely to have been overlooked elsewhere given its small size.
 - *Litosanthes biflora* is a new record for China mainland and had only been recorded in Hainan, Taiwan and the Philippines.
 - *Bulbophyllum* sp.1 may be new to science. Further morphological study is needed to confirm its identity.
 - *Eria rosea* has only been recorded in Hong Kong and Hainan.
 - *Eria thao* has only been recorded in Hainan and Vietnam.
 - *Dendrobim terminale* has only been recorded in southern Yunnan within China.
 - *Liparis cespitosa* has only been recorded in Yunnan and Taiwan within China.
- Among the plant species recorded, there are a number of species of conservation importance.
 - The orchids *Anoectochilus roxburghii*, *Cymbidium ensifolium* and *C. goeringii* are endangered in China due to over-collection, for medicinal and ornamental purposes.
 - *Hopea chinensis* is considered Critically Endangered globally and is also under Class I National Protection. It is endemic to South Guangxi but it is also likely to occur in North Vietnam. Large populations with abundant fruiting trees and saplings were found in the present surveys. One single large tree was also seen.
 - *Madhuca pasquieri* is Vulnerable globally and is under Class II National Protection. It is often the dominant species in northern subtropical lowland evergreen forest of South China. It was found to be locally abundant.

- *Diplopanax stachyanthus* is Vulnerable globally and is under Class II National Protection. A large population with more than 10 reproductive trees was found in evergreen forest at high altitude.
 - *Ixonanthes chinensis* is considered Vulnerable globally although it is abundant in northern tropical evergreen forest of South China.
 - *Semiliquidambar cathayensis* is under Class II National Protection and is considered to be Near-Threatened globally.
 - *Gymnosphaera hancockii* and *G. podophylla* belong to the tree fern family, which is under Class II National Protection. Both species have a widespread distribution in South China but the latter is common in most forest whereas the former is rare and restricted to the well-protected forest. A small population of *G. hancockii* with about 8-10 plants was seen.
 - *Brainea insignis* and *Cibotium barometz* are under Class II National Protection in China, although both species are widespread and abundant in relatively degraded landscape in South China.
 - *Phlegmariurus shangsiensis* is endemic to South Guangxi. Only a few individuals were seen.
 - *Fissistigma cupreonitens* is endemic to South Guangxi. It is locally abundant at two locations, and a few individuals were seen at two other locations.
 - *Fissistigma shangzeense* is endemic to South Guangxi. A few individuals were seen.
 - *Castanopsis amabilis* is endemic to South Guangxi. It is found to be locally common and co-dominate in one of the lowland forests.
 - *Ardisia filiformis* is endemic to South Guangxi. It is locally common in one of the hillside forests at high altitude.
 - *Calamus austro-guangxiensis* is endemic to South Guangxi.
 - *Itea chingiana* is endemic to Guangxi but previous records were restricted to North Guangxi. Only a few individuals were seen.
 - *Lithocarpus attenuatus* is endemic to South Guangxi and South Guangdong. A few fruiting trees were seen in the 2000 survey.
 - *Ardisia pubivenula* is endemic to South Guangxi and Hainan. It is locally common one location.
 - *Olea tetragonoclada* is endemic to Guangxi.
 - *Ormosia sericeolucida* is endemic to South Guangxi and Southwest Guangdong. It is fairly common in the areas surveyed as a whole and was seen at a few other locations.
 - *Urophyllum chinense* is restricted to Southwest Guangdong, South Guangxi, Southeast Yunnan, and North Vietnam, and is only found in relatively well-preserved lowland forest. It is locally common at two locations.
 - *Xanthophytum kwangtungense* is restricted to South Guangxi, South Yunnan and North Vietnam. It is locally common at one location.
 - *Rhododendron emarginatum* is rare in China and had been recorded from South Guangxi, Guizhou, Southeast Yunnan and Vietnam. A small population was found.
 - *Horsfieldia glabra* is rare in China and is restricted to forest in tropical region (South Yunnan, South Guangxi and South Guangdong). It is locally common at one location.
 - The population of *Vanilla* sp. recorded in the 2000 survey is the largest wild population found in KFBG surveys.
 - All orchid species recorded are listed in CITES Appendix II.
- Of the 55 orchid species recorded, 20 (36%) were terrestrial and 35 (64%) were epiphytic. The occurrence of tropical genera (e.g. *Vanilla*) and the high proportion of epiphytic orchids indicate Shiwandashan's forest has tropical characteristics.

Table 1. Vascular plants of Shiwandashan National Nature Reserve recorded in the present surveys (excluding Orchidaceae)t. Species which are nationally Protected (Class I or II) (State Forestry Administration & Ministry of Agriculture, 1999), globally Threatened or Lower Risk (Near-threatened) (IUCN, 2002) or globally restricted are indicated.

Family	Scientific name	Sep 2000	Dec 2001	Remarks
Pteridophyta				
Adiantaceae	<i>Adiantum flabellulatum</i> L.		✓	
Aspidiaceae	<i>Hemigramma decurrens</i> (Hook.) Copel.		✓	
	<i>Pleocnemia wintii</i> Holtt.		✓	
	<i>Tectaria phaeocaulis</i> (Rosenst.) C. Chr.		✓	

Family	Scientific name	Sep 2000	Dec 2001	Remarks
Aspleniaceae	<i>Asplenium neolaserpitiifolium</i> Tardieu & Ching		✓	
	<i>Asplenium normale</i> D. Don		✓	
	<i>Asplenium prolongatum</i> Hook.		✓	
	<i>Neottopteris nidus</i> (L.) J. Sm.		✓	
	<i>Neottopteris phyllitidis</i> (D. Don) J. Sm.		✓	
Athyriaceae	<i>Diplazium donianum</i> (Mett.) Tardieu		✓	
Blechnaceae	<i>Blechnum orientale</i> L.		✓	
	<i>Brainea insignis</i> (Hook.) J. Sm.		✓	Protected II
	<i>Chieniopteris harlandii</i> (Hook.) Ching		✓	
	<i>Woodwardia japonica</i> (L.f.) Sm.		✓	
	<i>Woodwardia orientalis</i> Sw.	✓	✓	
Bolbitidaceae	<i>Bolbitis heteroclita</i> (C. Presl) Ching		✓	
	<i>Bolbitis subcordata</i> (Copel.) Ching		✓	
Cyatheaceae	<i>Gymnosphaera hancockii</i> (Copel.) Ching		✓	Protected II
	<i>Gymnosphaera podophylla</i> (Hook.) Copel.	✓	✓	Protected II
Davalliaceae	<i>Humata repens</i> (L.f.) Diels		✓	
Dennstaedtiaceae	<i>Microlepia hookeriana</i> (Wall. ex Hook.) C. Presl.		✓	
Dicksoniaceae	<i>Cibotium barometz</i> (L.) J. Sm.		✓	Protected II
Dryopteridaceae	<i>Arachniodes sphaerosora</i> (Ching) Ching	✓	✓	
Elaphoglossaceae	<i>Elaphoglossum yoshinagae</i> (Yatabe) Makino		✓	
Gleicheniaceae	<i>Dicranopteris pedata</i> (Houtt.) Nakaike		✓	
	<i>Diplopterygium chinensis</i> (Rosenst.) DeVol		✓	
Grammitidaceae	<i>Grammitis dorsipila</i> (Christ) C. Chr. & Tardieu		✓	
	<i>Prosaptia khasyana</i> (Hook.) C. Chr. & Tardieu	✓	✓	
Huperziaceae	<i>Huperzia serrata</i> (Thunb.) Trevis.		✓	
	<i>Phlegmariurus shangsiensis</i> C. Y. Yang		✓	endemic to S. Guangxi
Hymenophyllaceae	<i>Mecodium badium</i> (Hook. & Grev.) Ching		✓	
	<i>Mecodium excertum</i> (Wall.) Copel.		✓	new record for Guangxi
	<i>Mecodium microsorum</i> (Bosch) Ching		✓	
	<i>Mecodium osmundoides</i> (Bosch) Ching		✓	
	<i>Meringium denticulatum</i> (Sw.) Copel.		✓	
	<i>Microgonium beccarianum</i> (Cesati) Copel		✓	new record for China mainland
Lindsaeaceae	<i>Lindsaea heterophylla</i> Dryand.		✓	
	<i>Lindsaea orbiculata</i> (Lam.) Mett. ex Kuhn		✓	
	<i>Stenoloma chusanum</i> (L.) Ching		✓	
Loxogrammaceae	<i>Loxogramme salicifolia</i> (Makino) Makino		✓	
Lycopodiaceae	<i>Lycopodium casuarinoides</i> (Spring) Holub		✓	
	<i>Palhinhaea cernua</i> (L.) Franco & Vasc.		✓	
Lygodiaceae	<i>Lygodium japonicum</i> (Thunb.) Sw.	✓		
Marattiaceae	<i>Angiopteris</i> sp.		✓	
Nephrolepidiaceae	<i>Nephrolepis auriculata</i> (L.) Trimea		✓	
Osmundaceae	<i>Osmunda vachellii</i> Hook.	✓	✓	
Polypodiaceae	<i>Colysis elliptica</i> (Thunb.) Ching var. <i>pothifolia</i> Ching		✓	
	<i>Colysis wrightii</i> (Hook.) Ching		✓	
	<i>Lepidogrammits rostrata</i> (Bedd.) Ching		✓	
	<i>Lepidomicrosorium buergerianum</i> (Miq.) Bosman		✓	
	<i>Microsorium fortunei</i> (T. Moore) Ching		✓	
	<i>Pyrrosis lingua</i> (Thunb.) Farw		✓	
	<i>Pteris incisa</i> (Thunb.) J. Sm.		✓	
Pteridaceae	<i>Pteris semipinnata</i> L.		✓	
	<i>Pteridium aquilinum</i> (L.) Kuhn var. <i>latiusculum</i> (Desv.) Underw. ex A. Heller		✓	
	<i>Selaginella delicatula</i> (Desv. ex Poir.) Alston		✓	
Selaginellaceae	<i>Selaginella delicatula</i> (Desv. ex Poir.) Alston		✓	
Thelypteridaceae	<i>Pronephrium simplex</i> (Hook.) Holttum		✓	
Thelypteridaceae	<i>Pronephrium triphyllum</i> (Sw.) Holttum		✓	
Vittariaceae	<i>Vittaria flexuosa</i> Fée		✓	
Gymnospermae				
Gnetaceae	<i>Gnetum luofuense</i> C. Y. Cheng		✓	
	<i>Gnetum montanum</i> Markgr.	✓		
	<i>Gnetum parvifolium</i> (Warb.) Chun		✓	
Pinaceae	<i>Pinus massoniana</i> Lamb.	✓	✓	

Family	Scientific name	Sep 2000	Dec 2001	Remarks
Podocarpaceae	<i>Podocarpus neriifolius</i> D. Don	✓	✓	
Taxodiaceae	<i>Cunninghamia lanceolata</i> (Lamb.) Hook.		✓	cultivated
Angiospermae				
Dicotyledoneae				
Aceraceae	<i>Acer davidii</i> Franch.		✓	
Actinidiaceae	<i>Actinidia glaucophylla</i> F. Chun		✓	
	<i>Actinidia latifolia</i> (Gardner & Champ.) Merr.	✓	✓	
	<i>Actinidia liangguangensis</i> C.F. Liang	✓		
	<i>Saurauia tristyla</i> DC.	✓	✓	
Anacardiaceae	<i>Choerospondias axillaris</i> (Roxb.) B.L. Burt & A.W. Hill		✓	
	<i>Rhus chinensis</i> Mill.	✓	✓	
	<i>Toxicodendron succedaneum</i> (L.) Kuntze.	✓	✓	
Annonaceae	<i>Artabotrys hongkongensis</i> Hance		✓	
	<i>Desmos chinensis</i> Lour.		✓	
	<i>Fissistigma cupreonitens</i> Merr. & Chun	✓	✓	endemic to S. Guangxi
	<i>Fissistigma glaucescens</i> (Hance) Merr.		✓	
	<i>Fissistigma maclurei</i> Merr.		✓	
	<i>Fissistigma oldhamii</i> (Hemsl.) Merr.		✓	
	<i>Fissistigma shangtzeense</i> Tsiang & P.T. Li	✓	✓	endemic to S. Guangxi
	<i>Goniothalamus chinensis</i> Merr. et Chun	✓		
	<i>Uvaria boniana</i> Finet & Gagnep.		✓	
	<i>Uvaria microcarpa</i> Champ. ex Benth.	✓	✓	
	<i>Uvaria tonkinensis</i> Finet & Gagnep.		✓	
Apocynaceae	<i>Chonemorpha eriostylis</i> Pit.		✓	
	<i>Melodinus suaveolens</i> Champ. ex Benth.		✓	
Aquifoliaceae	<i>Ilex ficoidea</i> Hemsl.		✓	
	<i>Ilex hainanensis</i> Merr.	✓		
	<i>Ilex rotunda</i> Thunb.	✓		
	<i>Ilex viridis</i> Champ. ex Benth.		✓	
Araliaceae	<i>Aralia armata</i> (Wall.) Seem.	✓	✓	
	<i>Dendropanax hainanensis</i> (Merr. & Chun) Merr. & Chun		✓	
	<i>Dendropanax proteus</i> Benth.		✓	
	<i>Diplopanax stachyanthus</i> Hand.-Mazz.		✓	Vulnerable, Protected II
	<i>Heteropanax fragrans</i> (D. Don) Seem.	✓	✓	
	<i>Schefflera leucantha</i> R. Vig.		✓	
	<i>Schefflera octophylla</i> (Lour.) Harms	✓	✓	
Asclepiadaceae	<i>Graphistemma pictum</i> (Champ. ex Benth.) Benth. & Hook. f. ex Maxim.		✓	
	<i>Hoya fusca</i> Wall.		✓	
	<i>Streptocaulon juvenas</i> (Lour.) Merr.	✓	✓	
Asteraceae	<i>Ageratum houstonianum</i> Mill.		✓	exotics from S. America
	<i>Ainsliaea trinervis</i> Y.Q. Tseng	✓		
	<i>Blumea riparia</i> DC.		✓	
	<i>Senecio scandens</i> Buch.-Ham.		✓	
	<i>Vernonia arborea</i> Buch.-Ham.		✓	
	<i>Vernonia cumingiana</i> Benth.		✓	
	<i>Vernonia solanifolia</i> Benth.		✓	
Balanophoraceae	<i>Balanophora spicata</i> Hayata		✓	
Begoniaceae	<i>Begonia crassirostris</i> Irmsch.		✓	
	<i>Begonia palmata</i> D. Don		✓	
Burseraceae	<i>Canarium album</i> (Lour.) Raeusch.	✓	✓	
Caesalpinaceae	<i>Caesalpinia crista</i> L.	✓	✓	
	<i>Pterolobium punctatum</i> Hemsl.	✓		
Campanulaceae	<i>Pentaphragma spicatum</i> Merr.	✓		
Caprifoliaceae	<i>Viburnum fordiae</i> Hance		✓	
	<i>Viburnum odoratissimum</i> Ker Gawl.		✓	
Celastraceae	<i>Celastrus monospermus</i> Roxb.	✓	✓	
	<i>Euonymus hederaceus</i> Champ. ex Benth.		✓	
	<i>Euonymus mitratus</i> Pierre		✓	new record for Guangxi

Family	Scientific name	Sep 2000	Dec 2001	Remarks
Chloranthaceae	<i>Sarcandra glabra</i> (Thunb.) Nakai		✓	
Clethraceae	<i>Clethra faberi</i> Hance		✓	
Clusiaceae	<i>Calophyllum membranaceum</i> Gardner & Champ.		✓	
	<i>Cratoxylum cochinchinense</i> (Lour.) Blume		✓	
	<i>Cratoxylum formosum</i> (Jack) Dyer	✓		
	<i>Cratoxylum formosum</i> (Jack) Dyer subsp. <i>pruniflorum</i> (Kurz) Gogelin	✓		
	<i>Garcinia multiflora</i> Champ. ex Benth.	✓	✓	
Connaraceae	<i>Garcinia oblongifolia</i> Champ. ex Benth.		✓	
	<i>Rourea microphylla</i> (Hook. & Arn.) Planch.		✓	
Convolvulaceae	<i>Merremia umbellata</i> (L.) Hallier. f.		✓	
Cornaceae	<i>Aucuba chinensis</i> Benth.		✓	
Cucurbitaceae	<i>Trichosanthes pedata</i> Merr. & Chun	✓		
Daphniphyllaceae	<i>Daphniphyllum calycinum</i> Benth	✓	✓	
	<i>Daphniphyllum oldhami</i> (Hemsl.) Rosenth.	✓	✓	
	<i>Daphniphyllum</i> sp.	✓	✓	
Dilleniaceae	<i>Dillenia turbinata</i> Finet & Gagnep.	✓	✓	
	<i>Tetracera asiatica</i> (Lour.) Hoog.	✓	✓	
Dipterocarpaceae	<i>Hopea chinensis</i> Hand.-Mazz.	✓	✓	endemic to S. Guangxi, Critically Endangered, Protected I
Ebenaceae	<i>Diospyros morrisiana</i> Hance ex. Walpers	✓	✓	
Elaeocarpaceae	<i>Elaeocarpus chinensis</i> (Gardner & Champ.) Hook. f. ex Benth.		✓	
	<i>Elaeocarpus nitentifolius</i> Merr. & Chun		✓	
	<i>Elaeocarpus sylvestris</i> (Lour.) Poir.	✓	✓	
	<i>Elaeocarpus varunua</i> Buch.-Ham.	✓		
	<i>Sloanea sinensis</i> (Hance) Hemsl.	✓		
Ericaceae	<i>Craibiodendron kwangtungense</i> S. Y. Hu		✓	
	<i>Enkianthus quinqueflorus</i> Lour.		✓	
	<i>Gaultheria leucocarpa</i> Blume var. <i>crenulata</i> (Kurz) T.Z. Hsu	✓		
	<i>Lyonia ovalifolia</i> (Wall.) Drude		✓	
	<i>Rhododendron emarginatum</i> Hemsl. & E.H. Wilson		✓	
	<i>Rhododendron haofui</i> Chun & W.P. Fang		✓	
	<i>Rhododendron moullainense</i> Hook. f.		✓	
	<i>Rhododendron simsii</i> Planch.		✓	
<i>Vaccinium chunii</i> Merr. ex Sleumer			new record of Guangxi	
Erythroxylaceae	<i>Erythroxylum sinense</i> Y. C. Wu			
Escalloniaceae	<i>Itea chinensis</i> Hook. & Arn	✓	✓	
	<i>Itea chingiana</i> S.Y. Jin		✓	endemic to Guangxi
Euphorbiaceae	<i>Alchornea trewioides</i> (Benth.) Muell.-Arg.		✓	
	<i>Antidesma fordii</i> Hemsl.	✓	✓	
	<i>Antidesma japonicum</i> Siebold & Zucc.	✓		
	<i>Aporosa dioica</i> (Roxb.) Müll. Arg.		✓	
	<i>Bischofia javanica</i> Blume		✓	
	<i>Breynia fruticosa</i> (L.) Hook. f.		✓	
	<i>Bridelia insulana</i> Hance (<i>B. balansae</i> Tutch.)		✓	
	<i>Endospermum chinense</i> Benth.	✓	✓	
	<i>Glochidion eriocarpum</i> Champ. ex Benth.		✓	
	<i>Glochidion wrightii</i> Benth.		✓	
	<i>Macaranga adenantha</i> Gagnep.	✓		
	<i>Macaranga auriculata</i> (Merr.) Airy Shaw		✓	
	<i>Macaranga sampsoni</i> Hance	✓	✓	
	<i>Macaranga henryi</i> (Pax & K. Hoffm.) Rehder	✓	✓	
	<i>Macaranga kurzii</i> (Kuntze) Pax & K. Hoffm.		✓	
	<i>Mallotus hookerianus</i> (Seem.) Müll. Arg.		✓	
	<i>Mallotus oblongifolius</i> (Miq.) Müll. Arg.		✓	
	<i>Mallotus paniculatus</i> (Lam.) Müll.-Arg.	✓	✓	
	<i>Mallotus philippinensis</i> (Lam.) Mull. Arg.		✓	
	<i>Microdesmis caseariifolia</i> Planch.		✓	
<i>Sapium discolor</i> (Champ. ex Benth.) Müll.-Arg.	✓	✓		

Family	Scientific name	Sep 2000	Dec 2001	Remarks
Fagaceae	<i>Vernicia montana</i> Lour.		✓	
	<i>Castanopsis amabilis</i> W.C. Cheng & C.S. Chao		✓	endemic to S. Guangxi
	<i>Castanopsis carlesii</i> (Hemsl.) Hayata	✓	✓	
	<i>Castanopsis eyrei</i> (Champ. ex Benth.) Tutcher		✓	
	<i>Castanopsis fissa</i> (Champ. ex Benth.) Rehder & E. H. Wilson	✓	✓	
	<i>Castanopsis hystrix</i> Miq.		✓	
	<i>Castanopsis tibetana</i> Hance		✓	
	<i>Cyclobalanopsis delicatula</i> (Chun & Tsiang) Y.C. Hsu & H. Wei Jen		✓	
	<i>Cyclobalanopsis neglecta</i> Schottky		✓	
	<i>Cyclobalanopsis patelliformis</i> (Chun) Y.C. Hsu & H.W. Jen	✓		
	<i>Lithocarpus attenuatus</i> (Skan) Rehder	✓		endemic to S. Guangxi & S. Guangdong
	<i>Lithocarpus corneus</i> (Lour.) Rehder	✓		
	<i>Lithocarpus cyrtocarpus</i> (Drake) A. Camus	✓	✓	
	<i>Lithocarpus elizabethae</i> (Tutcher) Rehder	✓	✓	
	<i>Lithocarpus fenestratus</i> (Roxb.) Rehder			
	<i>Lithocarpus hancei</i> (Benth.) Rehder	✓		
	<i>Lithocarpus longipedicellatus</i> (Hickel & A. Camus) A. Camus	✓		
	<i>Lithocarpus uvariifolius</i> (Hance) Rehder		✓	
	Flacourtiaceae	<i>Bennettiodendron brevipes</i> Merr.	✓	
<i>Bennettiodendron leprosipes</i> (Clos) Merr.			✓	
<i>Casearia balansae</i> Gagnep.			✓	
Gentianaceae	<i>Canscora andrographioides</i> Griffith ex C.B. Clarke	✓	✓	
Gesneriaceae	<i>Aeschynanthus acuminatus</i> Wall. ex A. DC.		✓	
	<i>Lysionotus pauciflorus</i> Maxim.		✓	
Haloragidaceae	<i>Rhynchosyche ellipticum</i> (Wal. ex D. Dietr.) A. DC.	✓		
Hamamelidaceae	<i>Haloragis micrantha</i> (Thunb.) R. Br.		✓	
	<i>Altingia chinensis</i> (Champ. ex Benth.) Oliv. ex Hance	✓	✓	
	<i>Corylopsis multiflora</i> Hance		✓	
	<i>Distylium myricoides</i> Hemsl.	✓		
	<i>Exbucklandia tonkinensis</i> (Lecomte) Steenis	✓		
	<i>Liquidambar formosana</i> Hance		✓	
	<i>Mytilaria laosensis</i> Lecomte	✓	✓	
	<i>Semiliquidambar cathayensis</i> H. T. Chang		✓	Protected II, Lower risk (nt)
	<i>Dichroa febrifuga</i> Lour.		✓	
Hydrangeaceae	<i>Pileostegia viburnoides</i> Hook. f. & Thomson		✓	
	<i>Gomphandra tetrandra</i> (Wall.) Sleum.	✓	✓	
Icacinaceae	<i>Mappianthes iodooides</i> Hand.-Mazz.	✓	✓	
	<i>Illicium verum</i> Hook. f.		✓	cultivated
Ixonanthaceae	<i>Ixonanthes chinensis</i> Champ.	✓	✓	Vulnerable
Juglandaceae	<i>Engelhardtia roxburghiana</i> Wall.	✓	✓	
Lauraceae	<i>Actinodaphne pilosa</i> (Lour.) Merr.	✓	✓	
	<i>Beilschmiedia intermedia</i> C.K. Allen	✓		
	<i>Cinnamomum cassia</i> (L.) Presl		✓	cultivated
	<i>Cinnamomum porrectum</i> (Roxb.) Kosterm.	✓	✓	
	<i>Cinnamomum validinerve</i> Hance		✓	
	<i>Cryptocarya chingii</i> W.C. Cheng		✓	
	<i>Cryptocarya concinna</i> Hance		✓	
	<i>Cryptocarya densiflora</i> Blume	✓		
	<i>Lindera caudata</i> (Nees) Hook. f.	✓	✓	
	<i>Lindera kwangtungensis</i> (H. Liu) C.K. Allen	✓		
	<i>Lindera pulcherrima</i> (Nees) Benth. var. <i>hemsleyana</i> (Diels) H.B. Cui	✓	✓	
	<i>Litsea cubeba</i> (Lour.) Pers.	✓	✓	
	<i>Litsea elongata</i> (Nees) Benth. & Hook. f.		✓	
	<i>Litsea greenmaniana</i> C.K. Allen	✓	✓	
	<i>Litsea lancilimba</i> Merr.	✓	✓	
	<i>Litsea rotundifolia</i> Hemsl. var. <i>oblongifolia</i> (Nees) C. K. Allen		✓	

Family	Scientific name	Sep 2000	Dec 2001	Remarks
	<i>Litsea variabilis</i> Hemsl. fo. <i>chinensis</i> (C.K. Allen) Yen	✓	✓	
	C. Yang & P.H. Huang			
	<i>Litsea verticillata</i> Hance		✓	
	<i>Machilus decursinervis</i> Chun		✓	
	<i>Machilus salicina</i> Hance	✓	✓	
	<i>Machilus velutina</i> Champ. ex Benth.	✓	✓	
	<i>Machilus wangchiana</i> Chun	✓	✓	
	<i>Neolitsea chuii</i> Merr.	✓		
	<i>Neolitsea kwangsiensis</i> H. Liu		✓	
	<i>Neolitsea pulchella</i> (Meissn.) Merr	✓	✓	
Loganiaceae	<i>Gelsemium elegans</i> (Gardner & Champ.) Benth.		✓	
Loranthaceae	<i>Scurrula parasitica</i> L.		✓	
Magnoliaceae	<i>Magnolia championii</i> Benth.	✓		
	<i>Magnolia paenetauma</i> Dandy	✓		
	<i>Manglietia chingii</i> Dandy	✓		
	<i>Manglietia fordiana</i> Oliv.	✓		
	<i>Michelia foveolata</i> Merr. ex Dandy	✓	✓	
	<i>Michelia maudiae</i> Dunn			
Malvaceae	<i>Urena lobata</i> L.	✓	✓	pantropical weed
Melastomataceae	<i>Barthea barthei</i> (Hance ex Benth.) Krasser		✓	
	<i>Blastus cavaleriei</i> H. Lévl. & Vaniot	✓		
	<i>Blastus cochinchinensis</i> Lour.	✓	✓	
	<i>Medinilla septentrionalis</i> (W.W. Sm.) H.L. Li	✓		
	<i>Melastoma candidum</i> D. Don		✓	
	<i>Melastoma dodecandrum</i> Lour.		✓	
	<i>Melastoma sanguineum</i> Sims		✓	
	<i>Memecylon ligustrifolium</i> Champ. ex Benth.	✓	✓	
	<i>Sonerila rivularis</i> Cogn.	✓		
Meliaceae	<i>Dysoxylum hongkongense</i> (Tutcher) Merr.	✓		
	<i>Trichilia connaroides</i> (Wight & Arn.) Benth. var. <i>microcarpa</i> (Pierre) Benth.		✓	
	<i>Trichilia sinensis</i> Benth.		✓	
Menispermaceae	<i>Albertisia laurifolia</i> Yamamoto	✓		
	<i>Hypserpa nitida</i> Miers		✓	
Mimosaceae	<i>Acacia pennata</i> (L.) Willd.		✓	
	<i>Albizia corniculata</i> (Lour.) Druce	✓	✓	
	<i>Cylindrokelupha kerrii</i> (Gagnep.) T.L. Wu	✓	✓	
	<i>Cylindrokelupha tonkinensis</i> (I.C. Nielsen) T.L. Wu		✓	
	<i>Cylindrokelupha turgida</i> (Merr.) T.L. Wu	✓		
	<i>Entada phaseoloides</i> (L.) Merr.	✓	✓	
	<i>Pithecellobium clypearia</i> (Jack) Benth.		✓	
	<i>Pithecellobium lucidium</i> Benth.		✓	
Moraceae	<i>Artocarpus styracifolius</i> Pierre		✓	
	<i>Ficus abelii</i> Miq.		✓	
	<i>Ficus auriculata</i> Lour.	✓		
	<i>Ficus esquiroliana</i> H. Lévl.	✓	✓	
	<i>Ficus fistulosa</i> Reinw. ex Blume	✓	✓	
	<i>Ficus hirta</i> Vahl		✓	
	<i>Ficus hispida</i> L. f.	✓		
	<i>Ficus pyriformis</i> Hook. & Arn.		✓	
	<i>Ficus superba</i> (Miq.) Miq.		✓	
	<i>Ficus variolosa</i> Lindl. ex Benth.	✓	✓	
	<i>Streblus indica</i> (Bureau) Corner	✓	✓	
Myricaceae	<i>Myrica rubra</i> (Lour.) Sieb. & Zucc.		✓	
Myristicaceae	<i>Horsfieldia glabra</i> (Reinw. ex Blume) Warb.		✓	
Myrsinaceae	<i>Ardisia filiformis</i> E. Walker		✓	endemic to S. Guangxi
	<i>Ardisia hanceana</i> Mez	✓	✓	
	<i>Ardisia mamillata</i> Hance		✓	
	<i>Ardisia pubivenula</i> E. Walker	✓		endemic to S. Guangxi & Hainan
	<i>Ardisia quinquegona</i> Blume	✓	✓	
	<i>Ardisia villosa</i> Roxb.		✓	
	<i>Ardisia waitakii</i> C.M. Hu			

Family	Scientific name	Sep 2000	Dec 2001	Remarks
	<i>Embelia laeta</i> (L.) Mez		✓	
	<i>Embelia parviflora</i> Wall. ex A. DC.		✓	
	<i>Embelia ribes</i> Burm. f.		✓	
	<i>Maesa acuminatissima</i> Merr.		✓	
	<i>Maesa japonica</i> (Thunb.) Moritz & Zoll.	✓	✓	
	<i>Maesa perlarius</i> (Lour.) Merr.		✓	
	<i>Mysine kwangsiensis</i> (E. Walker) Pipoly & C. Chen	✓		
	<i>Mysine seguinii</i> H. Lév		✓	
Myrtaceae	<i>Acmena acuminatissima</i> (Blume) Merr. & L. M. Perry	✓	✓	
	<i>Baeckea frutescens</i> L.		✓	
	<i>Eucalyptus citriodora</i> Hook. f.		✓	cultivated
	<i>Rhodomyrtus tomentosa</i> (Aiton) Hassk.	✓	✓	
	<i>Syzygium araiocladum</i> Merr. & L.M. Perry		✓	
	<i>Syzygium brachyantherum</i> Merr. & L.M. Perry	✓		
	<i>Syzygium buxifolium</i> Hook. & Arn.		✓	
	<i>Syzygium fluviatile</i> (Hemsl.) Merr. & L.M. Perry	✓	✓	
	<i>Syzygium hancei</i> Merr. & L. M. Perry		✓	
	<i>Syzygium imitans</i> Merr. & L.M. Perry	✓	✓	
	<i>Syzygium rehderianum</i> Merr. & L.M. Perry	✓		
Oleaceae	<i>Chionanthus ramiflorus</i> Roxb.		✓	
	<i>Jasminum lanceolarium</i> Roxb.		✓	
	<i>Olea tetragonoclada</i> L.C. Chia	✓		endemic to Guangxi
	<i>Osmanthus matsumuranus</i> Hayata	✓	✓	
Opiliaceae	<i>Urobotrya latisquama</i> (Gagnep.) Hiepko		✓	
Papilionaceae	<i>Bowringia callicarpa</i> Champ. ex Benth.		✓	
	<i>Dalbergia hancei</i> Benth.	✓	✓	
	<i>Dalbergia pinnata</i> (Lour.) Prain	✓		
	<i>Fordia cauliflora</i> Hemsl.	✓		
	<i>Millettia pachyloba</i> Drake		✓	
	<i>Ormosia merrilliana</i> L. Chen	✓		
	<i>Ormosia sericeolucida</i> L. Chen		✓	endemic to S. Guangxi & SW Guangdong
Pentaphylacaceae	<i>Pentaphylax euryoides</i> Gardner & Champ.		✓	
Piperaceae	<i>Piper hongkongense</i> C. DC.		✓	
Pittosporaceae	<i>Pittosporum balansae</i> Aug. DC.		✓	
	<i>Pittosporum balansae</i> Aug. DC. var. <i>angustifolium</i> Gagnep.		✓	
Polygalaceae	<i>Xanthophyllum hainanense</i> Hu	✓	✓	
Proteaceae	<i>Helicia cochinchinensis</i> Lour.		✓	
	<i>Helicia hainanensis</i> Hayata		✓	
	<i>Helicia longipetiolata</i> Merr. & Chun		✓	
	<i>Heliciopsis terminalis</i> (Kurz) Sleumer	✓	✓	
Ranunculaceae	<i>Clematis crassifolia</i> Benth.		✓	
Rhizophoraceae	<i>Carallia longipes</i> Chun ex W.C. Ko		✓	
Rosaceae	<i>Eriobotrya fragrans</i> Champ. ex Benth.		✓	
	<i>Laurocerasus undulata</i> (Buch.-Ham. ex D. Don) Roem.		✓	
	<i>Pygeum topengii</i> Merr.		✓	
	<i>Rhaphiolepis indica</i> (L.) Lindl.	✓	✓	
	<i>Rhaphiolepis lanceolata</i> Hu		✓	
	<i>Rhaphiolepis salicifolia</i> Lindl.		✓	
	<i>Rubus leucanthus</i> Hance		✓	
	<i>Rubus reflexus</i> Ker		✓	
	<i>Sorbus caloneura</i> (Stapf) Rehder	✓		
Rubiaceae	<i>Adina pilulifera</i> (Lam.) Franch. ex Drake		✓	
	<i>Canthium dicoccum</i> (Gaertn.) Teysmann & Binnedijk		✓	
	<i>Damnacanthus labordei</i> (H. Lév.) H.S. Lo		✓	
	<i>Diplospora dubia</i> (Lindl.) Masam.		✓	
	<i>Gardenia jasminoides</i> J. Ellis		✓	
	<i>Gardenia stenophylla</i> Merr.	✓	✓	
	<i>Hedyotis hedyotideia</i> (DC.) Merr.		✓	
	<i>Hedyotis platystipula</i> Merr.	✓	✓	
	<i>Ixora henryi</i> H. Lév.		✓	

Family	Scientific name	Sep 2000	Dec 2001	Remarks
	<i>Lasianthus chinensis</i> (Champ. ex Benth.) Benth.	✓	✓	
	<i>Lasianthus curtisii</i> King & Gamble		✓	
	<i>Lasianthus formosensis</i> Matsum.		✓	
	<i>Lasianthus sikkimensis</i> Hook. f.	✓	✓	
	<i>Lasianthus wallichii</i> (Wight & Arn.) Wight		✓	
	<i>Litosanthes biflora</i> Blume		✓	new record of China mainland
	<i>Metadina trichotoma</i> (Zoll. & Moritzi) Bakh. f.	✓		
	<i>Mussaenda erosa</i> Champ. ex Benth.		✓	
	<i>Pavetta hongkongensis</i> Brem.		✓	
	<i>Psychotria asiatica</i> L.	✓	✓	
	<i>Psychotria serpens</i> L.		✓	
	<i>Psychotria tutcheri</i> Dunn		✓	
	<i>Tarenna mollissima</i> (Hook. & Arn.) B.L. Rob.		✓	
	<i>Uncaria rhynchophylla</i> (Miq.) Miq. ex Havil.		✓	
	<i>Uncaria rhynchophylloides</i> F.C. How	✓	✓	
	<i>Urophyllum chinense</i> Merr. & Chun		✓	restricted to SW Guangdong, S Guangxi, S Yunnan & N Vietnam
	<i>Wendlandia formosana</i> Cowan subsp. <i>breviflora</i> F.C. How		✓	
	<i>Wendlandia uvariifolia</i> Hance	✓	✓	
	<i>Xanthophytum kwangtungense</i> (Chun & F.C. How) H.S. Lo	✓	✓	restricted to S Guangxi, S Yunnan & N Vietnam
Rutaceae	<i>Acronychia pedunculata</i> (L.) Miq.		✓	
	<i>Evodia leptota</i> (Spreng.) Merr.	✓	✓	
	<i>Skimmia arborescens</i> T. Anderson ex Gamble		✓	
Sabiaceae	<i>Meliosma angustifolia</i> Merr.		✓	
	<i>Meliosma rigida</i> Siebold & Zucc.	✓	✓	
	<i>Meliosma squamulata</i> Hance		✓	
	<i>Meliosma thorelii</i> Lecomte		✓	
	<i>Sabia limoniacea</i> Wall. ex Hook. f. & Thomson		✓	
Santalaceae	<i>Dendrotrophe frutescens</i> (Champ. ex Benth.) Danser		✓	
Sapotaceae	<i>Eberhardtia aurata</i> (Pierre ex Dubard) Lecomte	✓	✓	
	<i>Madhuca pasquieri</i> (Dubard) H.J. Lam	✓	✓	Protected II, Vulnerable
	<i>Sarcosperma laurinum</i> (Benth.) Hook. f.	✓	✓	
	<i>Sinosideroxylon wightianum</i> (Hook. & Arn.) Aubrév.		✓	
Sterculiaceae	<i>Helicteres angustifolia</i> L.		✓	
	<i>Pterospermum heterophyllum</i> Hance		✓	
	<i>Reevesia thyrsoidea</i> Lindl	✓		
	<i>Sterculia lanceolata</i> Cav.	✓	✓	
Styracaceae	<i>Alniphyllum fortunei</i> (Hemsl.) Makino	✓	✓	
Symplocaceae	<i>Symplocos adenophylla</i> Wall. ex G. Don		✓	
	<i>Symplocos adenopus</i> Hance	✓		
	<i>Symplocos cochinchinensis</i> (Lour.) S. Moore		✓	
	<i>Symplocos lancifolia</i> Siebold & Zucc.		✓	
Theaceae	<i>Adinandra bockiana</i> E. Pritz var. <i>acutifolia</i> (Hand.-Mazz.) Kobuski	✓	✓	
	<i>Adinandra glischroloma</i> Hand.-Mazz. var. <i>jubata</i> (H.L. Li) Kobuski		✓	
	<i>Adinandra millettii</i> (Hook. & Arn.) Benth. & Hook. f. ex Hance		✓	
	<i>Adinandra nitida</i> Merr. ex H.L. Li		✓	
	<i>Camellia assimilis</i> Champ. ex Benth.		✓	
	<i>Camellia caudata</i> Wall.	✓	✓	
	<i>Camellia fluviatilis</i> Hand.-Mazz.		✓	
	<i>Cleyera pachyphylla</i> Chun ex H.T. Chang		✓	
	<i>Eurya groffii</i> Merr.		✓	
	<i>Eurya nitida</i> Korthals		✓	

Family	Scientific name	Sep 2000	Dec 2001	Remarks
	<i>Gordonia axillaris</i> (Roxb. ex Ker Gawl.) Dietr.		✓	
	<i>Hartia villosa</i> (Merr.) Merr.	✓	✓	
	<i>Schima bambusifolia</i> Hu	✓	✓	
	<i>Schima superba</i> Gardn. & Champ.	✓	✓	
	<i>Ternstroemia gymnanthera</i> (Wight & Arn.) Bedd.		✓	
Thymelaeaceae	<i>Daphne papyracea</i> Wall. ex Steud.		✓	
	<i>Wikstroemia nutans</i> Champ. ex Benth.		✓	
Ulmaceae	<i>Gironniera subaequalis</i> Planch.	✓	✓	
	<i>Trema cannabina</i> Lour. var. <i>dielsiana</i> (Hand.-Mazz.) C.J. Chan		✓	
	<i>Trema tomentosa</i> (Roxb.) Hara		✓	
Urticaceae	<i>Debregeasia longifolia</i> (Burm. f.) Wedd.	✓		
	<i>Debregeasia squamata</i> King ex Hook. f.	✓		
Verbenaceae	<i>Clerodendrum cyrtophyllum</i> Turcz.	✓	✓	
	<i>Clerodendrum fortunatum</i> L.		✓	
	<i>Vitex quinata</i> (Lour.) F.N. Williams			
Vitaceae	<i>Tetrastigma caudatum</i> Merr. & Chun	✓		
	<i>Tetrastigma planicaule</i> (Hook. f.) Gagnep.	✓	✓	
Monocotyledoneae				
Amaryllidaceae	<i>Curculigo capitulata</i> (Lour.) Kuntze		✓	
Araceae	<i>Acorus gramineus</i> Sol.	✓	✓	
	<i>Epipremnum pinnatum</i> (L.) Engl.	✓	✓	
	<i>Pothos chinensis</i> (Raf.) Merr.		✓	
	<i>Rhaphidophora hongkongensis</i> Schott		✓	
Areaceae	<i>Calamus austro-guangxiensis</i> S.J. Pei & S.Y.Chen	✓		endemic to S. Guangxi
	<i>Calamus tetradactylus</i> Hance	✓	✓	
	<i>Caryota monostachys</i> Becc.		✓	
	<i>Pinanga discolor</i> Burret	✓	✓	
Commelinaceae	<i>Amischotolype hispida</i> (Less. & A. Rich.) D.Y. Hong	✓	✓	
Cyperaceae	<i>Carex cruciata</i> Wahlenb.		✓	
	<i>Carex cryptostachys</i> Brongn.		✓	
	<i>Carex perakensis</i> C.B. Clarke		✓	
	<i>Carex scaposa</i> C.B. Clarke		✓	
	<i>Carex tenuispicula</i> T. Tang & S.Y.Liang		✓	new record of Guangxi
	<i>Carex truncatigluma</i> C.B. Clarke		✓	
	<i>Gahnia tristis</i> Nees		✓	
	<i>Hypolytrum nemorum</i> (Vahl) Spreng.		✓	
	<i>Mapania silhetensis</i> C.B. Clarke		✓	
	<i>Scleria harlandii</i> Hance		✓	
	<i>Scleria levis</i> Retz.		✓	
Liliaceae	<i>Dianella ensifolia</i> (L.) DC.		✓	
	<i>Ophiopogon platyphyllus</i> Merr. & Chun		✓	
	<i>Smilax lanceifolia</i> Roxb.		✓	
Marantaceae	<i>Phrynium rheedei</i> Suresh & Nicolson	✓	✓	
Musaceae	<i>Musa balbisiana</i> Colla		✓	
Pandanaceae	<i>Pandanus austrosinensis</i> T. L. Wu		✓	
	<i>Pandanus tectorius</i> Parkinson	✓		
Poaceae	<i>Aristida chinensis</i> Munro		✓	
	<i>Centotheca lappacea</i>		✓	
	<i>Ischaemum</i> sp.		✓	
	<i>Lophatherum gracile</i> Brongn.		✓	
	<i>Miscanthus floridulus</i> (Labill.) Warb. ex K. Schum & Lauterb.		✓	
	<i>Miscanthus sinensis</i> Andersson		✓	
	<i>Neyraudia arundinacea</i> (L.) Henr.		✓	
	<i>Themeda villosa</i> (Poir.) A. Camus		✓	
	<i>Thysanolaena maxima</i> (Roxb.) Kuntze		✓	
Stemonaceae	<i>Stemona tuberosa</i> Lour.		✓	
Zingiberaceae	<i>Alpinia chinensis</i> (J. König) Roscoe	✓	✓	
	<i>Alpinia kwangsiensis</i> T.L. Wu & S.J. Chen	✓	✓	
	<i>Alpinia strobiliformis</i> T. L. Wu & S. J. Chen		✓	
	<i>Hedychium villosum</i> Wall.	✓		

Table 2. Orchids recorded in Shiwandashan National Nature reserve and surrounding areas in September 2000 and December 2001.

Scientific name	Habitat	Remarks	Sep 2000	Dec 2000	Dec 2001
<i>Anoectochilus roxburghii</i> (Wall.) Lindl.	on forest floor with rich humus	terrestrial, Endangered	✓	✓	✓
<i>Anoectochilus</i> (cf. <i>elwesii</i>) sp.	on forest floor	terrestrial			✓
<i>Arundina graminifolia</i> (D. Don) Hochr.	on exposed grassy slope and beside stream	terrestrial	✓		✓
<i>Bulbophyllum affine</i> Lindl.	on rock	epiphytic			✓
<i>Bulbophyllum</i> (cf. <i>pentenveneris</i>) sp.	on rock in forest	epiphytic		✓	
<i>Bulbophyllum kwangtungense</i> Schltr.	on rock	epiphytic			✓
<i>Bulbophyllum levinei</i> Schltr.	on rock and on tree trunk	epiphytic			✓
<i>Bulbophyllum odoratissimum</i> (Sm.) Lindl.	on rock and on tree trunk beside stream	epiphytic	✓		
<i>Bulbophyllum</i> sp.1	mainly on rock, a few on tree trunk beside stream	epiphytic; may be new to science	✓		
<i>Bulbophyllum</i> sp.2	on rock in forest	epiphytic; may be new to China		✓	
<i>Bulbophyllum</i> sp.3	on rock beside waterfall	epiphytic		✓	
<i>Bulbophyllum</i> sp.4	on rock	epiphytic			✓
<i>Calanthe clavata</i> Lindl.	on forest floor	terrestrial			✓
<i>Calanthe</i> sp.	on floor beside stream	terrestrial	✓		
<i>Cephalantheropsis calanthoides</i> (Ames) T.S. Liu et H.J. Su	on forest floor	terrestrial			✓
<i>Cephalantheropsis</i> (cf. <i>gracilis</i>) sp.	on forest floor with rich humus	terrestrial	✓		
<i>Cheirostylis</i> sp.	on forest floor with rich humus	terrestrial			✓
<i>Cleisostoma paniculatum</i> (Ker Gawl.) Garay	on tree trunk in forest	epiphytic	✓		
<i>Coelogyne fimbriata</i> Lindl.	on rock	epiphytic			✓
<i>Collabium formosanum</i> Hayata	on rough rock surface and floor with rich humus	terrestrial			✓
<i>Collabium</i> (cf. <i>chinensis</i>) sp.	on floor beside stream	terrestrial			✓
<i>Cryptostylis arachnites</i> (Blume) Sw.	on forest floor	terrestrial			✓
<i>Cymbidium ensifolium</i> (L.) Sw.	on dense bamboo floor	terrestrial, Endangered			✓
<i>Cymbidium goeringii</i> (Rchb. f.) Rchb. f.	on dense bamboo floor	terrestrial, Endangered			✓
<i>Cymbidium floribundum</i> Lindl.	on rock with rich humus	epiphytic			✓
<i>Cymbidium</i> sp.	on tree trunk	epiphytic			✓
<i>Dendrobium densiflorum</i> Lindl.	on rock	epiphytic			✓
<i>Dendrobium longicornu</i> Lindl.	on rock in forest	epiphytic			✓
<i>Dendrobium terminale</i> Par. et Rchb. f.	on rock beside stream	epiphytic, new to Guangxi			✓
<i>Dendrobium</i> sp.1	on tree trunk	epiphytic			✓
<i>Dendrobium</i> sp.2	on rock	epiphytic			✓
<i>Epigeneium fargesii</i> (Finet) Gagnep.	on rock	epiphytic			✓
<i>Eria corneri</i> Rchb. f.	on rock	epiphytic			✓
<i>Eria pusilla</i> (Griff.) Lindl.	on rock and at base of tree trunk beside stream	epiphytic	✓		✓

Scientific name	Habitat	Remarks	Sep 2000	Dec 2000	Dec 2001
<i>Eria rosea</i> Lindl.	on rock	epiphytic, new to Guangxi			✓
<i>Eria thao</i> Gagnep.	on rock and on tree trunk	epiphytic, new to Guangxi	✓		✓
<i>Eria</i> sp.	on rock with rich humus	epiphytic, may be new to Guangxi			✓
<i>Gastrochilus</i> sp.	on tree trunk in forest	epiphytic	✓		
<i>Goodyera</i> (cf. <i>repens</i>) sp.	on forest floor	terrestrial, may be new to Guangxi			✓
<i>Habenaria tonkinensis</i> Seidenf.	in crevices with rich humus beside the stream	terrestrial	✓		
<i>Hetaeria cristata</i> Blume	on forest floor	terrestrial			✓
<i>Liparis balansae</i> Gagnep.	on rock	epiphytic			✓
<i>Liparis bootanensis</i> Griff.	on rock	epiphytic	✓		✓
<i>Liparis cespitosa</i> (Thou.) Lindl.	on rock in dense bamboo	epiphytic, new to Guangxi			✓
<i>Liparis nervosa</i> (Thunb) Lindl.	on forest floor	terrestrial	✓		✓
<i>Liparis</i> sp.1	on rock in forest	epiphytic			✓
<i>Liparis</i> sp.2	on rock	epiphytic			✓
<i>Mischobulbum cordifolium</i> (Hook. f.) Schltr.	on forest floor	terrestrial	✓		✓
<i>Ornithochilus difformis</i> (Lindl.) Schltr.	on tree trunk	epiphytic			✓
<i>Pholidota chinensis</i> Lindl.	on rock and on tree trunk	epiphytic	✓	✓	✓
<i>Platanthera minor</i> (Miq.) Rchb. F.	on slope beside the road	terrestrial	✓		
<i>Podochilus khasianus</i> Hook. f.	on rock beside the stream	epiphytic			
<i>Vanilla</i> sp.	on rock and on tree trunk in forest	epiphytic	✓		✓
unknown sp.	on tree trunk	epiphytic			✓

Mammals

- In May 1997 forest park staff were interviewed regarding the mammal fauna of Shiwandashan. Status of mammals is inferred based on their responses (Fellowes & Hau, 1997), records in Guangxi Forestry Survey and Planning Institute & Shiwandashan Nature Reserve (2002), and past distribution records from the Shiwandashan area (Wu, 1993; Zhang Y. *et al.*, 1997 and references therein), and is listed in Table 3.

Table 3. The inferred status of mammals at Shiwandashan National Nature Reserve, based on interviews with forest park staff (Fellows & Hau, 1997), on records in Guangxi Forestry Survey and Planning Institute & Shiwandashan Nature Reserve (2002), and on past distribution records, “SS” = “Shangsi”, “FC” = “Fangcheng”, “NM” = “Ningming”, “QZ” = “Qinzhou” (Wu, 1993; Zhang Y. *et al.*, 1997). “_” = “present”, “+” = “rare”, “++” = “common”, “+++” = “abundant”. Sequence follows D.E. Wilson & Cole (2000).

Scientific name	English name	Forest Park staff (May 1997)	Guangxi Forestry Survey (2002)	Wu (1993)	Zhang Y. <i>et al.</i> (1997)	Probable Status
<i>Crocidura attenuata</i>	Indochinese Shrew	(not asked)	—		NM, QZ	present
<i>Crocidura</i> (nr. <i>russula</i>) sp. (recorded as <i>C. russula</i>)	white-toothed shrew	(not asked)	—		NM, QZ	present
<i>Suncus murinus</i>	Asian House Shrew	(not asked)	—		SS, NM	present
<i>Chimarrogale himalayica</i>	Himalayan Water Shrew	(not asked)	—		NM	present
<i>Tupaia belangeri</i>	Northern Tree Shrew	-	—		SS, NM	present
<i>Rhinolophus affinis</i>	Intermediate Horseshoe Bat	(not asked)	—		NM	present
<i>Rhinolophus rouxii</i>	Rufous Horseshoe Bat	(not asked)	—			present
<i>Rhinolophus pusillus</i> (recorded as <i>R. blythi</i>)	Least Horseshoe Bat	(not asked)	—		NM	present
<i>Rhinolophus macrotis</i>	Big-eared Horseshoe Bat	(not asked)	—		NM	present
<i>Hipposideros bicolor</i>	Bicoloured Roundleaf Bat	(not asked)	—		NM	present
<i>Hipposideros pomona</i> (recorded as <i>H. p. sinensis</i>)	Pomona Roundleaf Bat	(not asked)	—			present
<i>Hipposideros pratti</i>	Pratt's Roundleaf Bat	(not asked)	—		NM	present
<i>Nyctalus noctula</i> (recorded as <i>N. velutinus</i>)	Noctule	(not asked)	—		SS, NM	present
<i>Pipistrellus abramus</i>	Japanese Pipistrelle	(not asked)	—			present
<i>Pipistrellus ceylonicus</i>	Kelaart's Pipistrelle	(not asked)	—		SS, NM	present
<i>Scotomanes heathi insularis</i>	Greater Asiatic Yellow Bat	(not asked)	—		SS	present
<i>Myotis mystacinus</i>	Whiskered Bat	(not asked)	—			present
<i>Nycticebus coucang</i>	Slow Loris	-		NM	NM	uncertain
<i>Macaca arctoides</i>	Stump-tailed Macaque		—	SS	SS, NM	present
<i>Macaca assamensis</i>	Assam Macaque			NM	SS, NM	uncertain
<i>Macaca mulatta</i>	Rhesus Monkey		—	SS, NM	SS, NM	present
<i>Trachypithecus francoisi</i>	Francois's Leaf Monkey			SS, NM	SS, NM	uncertain
<i>Canis lupus</i>	Grey Wolf			SS, NM	SS, NM	insecure/extirpated
<i>Cuon alpinus</i>	Dhole				SS, NM	insecure/extirpated
<i>Nyctereutes procyonoides</i>	Raccoon Dog	+	—		SS, NM	insecure
<i>Vulpes vulpes</i>	Red Fox		—		SS, NM	insecure
<i>Catopuma temminckii</i>	Asiatic Golden Cat	+	—	SS, FC, NM	SS, NM, FC	insecure
<i>Prionailurus bengalensis</i>	Leopard Cat	+	—		SS, NM	insecure
<i>Neofelis nebulosa</i>	Clouded Leopard	+	—	SS, NM	SS, NM	insecure

Scientific name	English name	Forest Park staff (May 1997)	Guangxi Forestry Survey (2002)	Wu (1993)	Zhang Y. <i>et al.</i> (1997)	Probable Status
<i>Panthera pardus</i>	Leopard		—	SS	SS, NM	extirpated
<i>Panthera tigris</i>	Tiger		—		SS, NM	extirpated
<i>Herpestes javanicus</i>	Small Asian Mongoose		—		SS, NM	present
<i>Herpestes urva</i>	Crab-eating Mongoose	+	—		SS, NM	insecure
<i>Amblonyx cinereus</i>	Oriental Small-clawed Otter	+	—		SS, NM, QZ	insecure
<i>Lutra lutra</i>	Eurasian Otter		—		SS, NM, QZ	insecure
<i>Arctonyx collaris</i>	Hog Badger		—		NM, QZ	insecure
<i>Martes flavigula</i>	Yellow-throated Marten		—		SS, NM	insecure
<i>Meles meles</i>	Eurasian Badger		—		NM, QZ	insecure
<i>Melogale moschata</i>	Chinese Ferret-badger		—		SS, NM	insecure
<i>Mustela kathiah</i>	Yellow-bellied Weasel	+	—		NM	present
<i>Mustela sibirica</i>	Siberian Weasel	+	—		SS, NM	present
<i>Ursus thibetanus</i>	Asiatic Black Bear		—		NM	insecure
<i>Chrotogale owstoni</i>	Owston's Banded Civet		—	NM	NM	uncertain
<i>Paguma larvata</i>	Masked Palm Civet	++	—		SS, NM	present
<i>Paradoxurus hermaphroditus</i>	Asian Palm Civet	++	—	SS, NM	SS, NM	present
<i>Prionodon pardicolor</i>	Spotted Linsang	++	—		SS, NM	present
<i>Viverra zibetha</i>	Large Indian Civet	+	—		SS, NM	insecure
<i>Viverricula indica</i>	Small Indian Civet	+	—		SS, NM	insecure
<i>Sus scrofa</i>	Wild Boar	++	—		SS, NM, QZ	present
<i>Moschus berezovskii</i>	Chinese Forest Musk Deer		—		SS, YC	insecure
<i>Cervus unicolor</i>	Sambar		—		SS, NM	insecure
<i>Muntiacus muntjak</i>	Indian Muntjac	++	—		SS, NM, QZ	present
<i>Muntiacus reevesii</i>	Reeves's Muntjac	++	—		SS, NM	present
<i>Naemorhedus caudatus</i>	Chinese Goral	+	—			insecure
<i>Naemorhedus sumatraensis</i>	Serow	+	—		SS, NM	insecure
<i>Manis pentadactyla</i>	Chinese Pangolin	+	—		SS, NM	insecure
<i>Callosciurus erythraeus</i>	Pallas's Squirrel	++	—		NM	present
<i>Dremomys pernyi</i>	Perny's Long-nosed Squirrel		—		NM	present
<i>Dremomys rufigenis</i>	Asian Red-cheeked Squirrel		—		SS, NM, FC	present
<i>Ratufa bicolor</i>	Black Giant Squirrel		—	FC	NM	insecure
<i>Tamias maritimus</i>	Maritime Striped Squirrel		—		SS	present
<i>Trogopterus pearsonii</i>	Hairy-footed Flying Squirrel		—	SS, NM	SS, NM	present

Scientific name	English name	Forest Park staff (May 1997)	Guangxi Forestry Survey (2002)	Wu (1993)	Zhang Y. et al. (1997)	Probable Status
<i>Petaurista alborufus</i>	Red and White Flying Squirrel		—		SS, NM	present
<i>Rhizomys pruinosus</i>	Hoary Bamboo Rat	++	—		SS, NM	present
<i>Rhizomys sinensis</i>	Chinese Bamboo Rat		—		SS, NM	present
<i>Hystrix brachyura</i>	Malayan Porcupine	+	—			present
<i>Atherurus macrourus</i>	Asiatic Brush-tailed Porcupine		—		SS, NM, FC	present
<i>Lepus capensis</i>	Brown Hare		—			uncertain
<i>Lepus sinensis</i>	Chinese Hare	++	—		SS, NM	present

- Some of the species suspected to occur are of particular conservation importance:
 - Clouded Leopard *Neofelis nebulosa* is Vulnerable globally and Class I Protected nationally.
 - Stump-tailed Macaque *Macaca arctoides*, Dhole *Cuon alpinus*, Asiatic Golden Cat *Catopuma temminckii*, Eurasian Otter *Lutra lutra*, Asiatic Black Bear *Ursus thibetanus* and Chinese Goral *Naemorhedus caudatus* are Vulnerable globally and Class II Protected nationally.
 - Malayan Porcupine *Hystrix brachyura* is Vulnerable globally.
 - Rhesus Monkey *Macaca mulatta*, Oriental Small-clawed Otter *Amblonyx cinereus*, Chinese Forest Musk Deer *Moschus berezovskii* and Chinese Pangolin *Manis pentadactyla* are at Lower Risk (Near-threatened) globally and Class II Protected nationally.
- Serow *Naemorhedus sumatraensis*, Yellow-throated Marten *Martes flavigula* and Black Giant Squirrel *Ratufa bicolor* are Class II Protected nationally.

Birds

- One hundred and eleven bird species were recorded in these surveys (Table 4).
- Among the most frequently encountered species in all three surveys were Light-vented Bulbul *Pycnonotus sinensis*, Chestnut Bulbul *Hypsipetes castanotus*, Black Bulbul *Hypsipetes leucocephalus*, streak-breasted Scimitar Babbler *Pomatorhinus ruficollis*, Rufous-capped Babbler *Stachyris ruficeps* and Grey-cheeked Fulvetta *Alcippe morrisonia*.
- The timing of the three surveys coincided with spring, autumn and winter migrations, explaining the high richness of raptors, warblers, flycatchers and thrushs.

Table 4. Birds recorded in Siwandashan National Nature Reserve and Siwandashan National Forest Park, March and September 2000 and December 2001. Sequence follows Clements (2000).

Scientific name	English name
<i>Ardeola bacchus</i>	Chinese Pond Heron
<i>Bubulcus ibis</i>	Cattle Egret
<i>Butorides striatus</i>	Little Heron
<i>Pernis ptilorhynchus</i>	Oriental Honey Buzzard
<i>Spilornis cheela</i>	Crested Serpent Eagle
<i>Milvus migrans</i>	Black Kite
<i>Accipiter trivirgatus</i>	Crested Goshawk
<i>Accipiter badius</i>	Shikra
<i>Accipiter gularis</i>	Japanese Sparrowhawk
<i>Accipiter virgatus</i>	Besra
<i>Falco subbuteo</i>	Eurasian Hobby
<i>Bambusicola thoracica</i>	Chinese Bamboo Partridge
<i>Lophura nycthemera</i>	Silver Pheasant
<i>Turnix suscitator</i>	Barred Buttonquail
<i>Amaurornis phoenicurus</i>	White-breasted Waterhen
<i>Streptopelia chinensis</i>	Spotted Dove
<i>Hierococcyx sparverioides</i>	Large Hawk Cuckoo
<i>Hierococcyx fugax</i>	Hodgson's Hawk Cuckoo
<i>Centropus sinensis</i>	Greater Coucal
<i>Centropus bengalensis</i>	Lesser Coucal
<i>Otus spilocephalus</i>	Mountain Scops Owl
<i>Otus bakkamoena</i>	Collared Scops Owl
<i>Otus sunia</i>	Oriental Scops Owl
<i>Glaucidium cuculoides</i>	Asian Barred Owlet
<i>Hirundapus cochinchinensis</i>	White-vented Needletail
<i>Alcedo atthis</i>	Common Kingfisher
<i>Halcyon smyrnensis</i>	White-throated Kingfisher
<i>Halcyon pileata</i>	Black-capped Kingfisher
<i>Merops philippinus</i>	Blue-tailed Bee-eater
<i>Megalaima virens</i>	Great Barbet
<i>Megalaima oorti</i>	Black-browed Barbet
<i>Megalaima asiatica</i>	Blue-throated Barbet
<i>Picumnus innominatus</i>	Speckled Piculet
<i>Sasia ochracea</i>	White-browed Piculet
<i>Hirundo rustica</i>	Barn Swallow
<i>Hirundo daurica</i>	Red-rumped Swallow
<i>Delichon dasypus</i>	Asian House Martin
<i>Motacilla alba</i>	White Wagtail
<i>Motacilla cinerea</i>	Grey Wagtail
<i>Anthus hodgsoni</i>	Olive-backed Pipit
<i>Pericrocotus flammeus</i>	Scarlet Minivet
<i>Pericrocotus solaris</i>	Grey-chinned Minivet
<i>Hemipus picatus</i>	Bar-winged Flycatcher-shrike
<i>Pycnonotus jocosus</i>	Red-whiskered Bulbul
<i>Pycnonotus sinensis</i>	Light-vented Bulbul
<i>Pycnonotus aurigaster</i>	Sooty-headed Bulbul
<i>Alophoixus pallidus</i>	Puff-throated Bulbul
<i>Hypsipetes castanotus</i>	Chestnut Bulbul
<i>Hypsipetes mcclllandii</i>	Mountain Bulbul
<i>Hypsipetes leucocephalus</i>	Black Bulbul

Scientific name	English name
<i>Myophonus caeruleus</i>	Blue Whistling Thrush
<i>Zoothera citrina</i>	Orange-headed Thrush
<i>Zoothera dauma</i>	Scaly Thrush
<i>Turdus hortulorum</i>	Grey-backed Thrush
<i>Turdus cardis</i>	Japanese Thrush
<i>Turdus pallidus</i>	Pale Thrush
<i>Turdus chrysolaus</i>	Brown-headed Thrush
<i>Brachypteryx leucophrys</i>	Lesser Shortwing
<i>Luscinia sibilans</i>	Rufous-tailed Robin
<i>Tarsiger cyanurus</i>	Orange-flanked Bush Robin
<i>Prinia atrogularis</i>	Hill Prinia
<i>Prinia rufescens</i>	Rufescent Prinia
<i>Prinia flaviventris</i>	Yellow-bellied Prinia
<i>Prinia inornata</i>	Plain Prinia
<i>Cettia fortipes</i>	Brownish-flanked Bush Warbler
<i>Bradypterus seebohmi</i>	Russet Bush Warbler
<i>Orthotomus cuculatus</i>	Mountain Tailorbird
<i>Orthotomus sutorius</i>	Common Tailorbird
<i>Phylloscopus fuscatus</i>	Dusky Warbler
<i>Phylloscopus proregulus</i>	Pallas's Leaf Warbler
<i>Phylloscopus inornatus</i>	Yellow-browed Warbler
<i>Phylloscopus tenellipes</i>	Pale-legged Leaf Warbler
<i>Phylloscopus reguloides</i>	Blyth's Leaf Warbler
<i>Phylloscopus ricketti</i>	Sulphur-breasted Warbler
<i>Seicercus burkii</i>	Golden-spectacled Warbler
<i>Abroscopus albogularis</i>	Rufous-faced Warbler
<i>Muscicapa dauurica</i>	Asian Brown Flycatcher
<i>Niltava davidi</i>	Fujian Niltava
<i>Cyornis hainana</i>	Hainan Blue Flycatcher
<i>Rhipidura albicollis</i>	White-throated Fantail
<i>Copsychus saularis</i>	Oriental Magpie Robin
<i>Phoenicurus aureus</i>	Daurian Redstart
<i>Rhyacornis fuliginosus</i>	Plumbeous Water Redstart
<i>Enicurus schistaceus</i>	Slaty-backed Forktail
<i>Garrulax maesi</i>	Grey Laughingthrush
<i>Garrulax canorus</i>	Hwamei
<i>Garrulax sannio</i>	White-browed Laughingthrush
<i>Pomatorhinus ruficollis</i>	Streak-breasted Scimitar Babbler
<i>Napothera brevicaudata</i>	Streaked Wren Babbler
<i>Pnoepyga pusilla</i>	Pygmy Wren Babbler
<i>Stachyris ruficeps</i>	Rufous-capped Babbler
<i>Leiothrix lutea</i>	Red-billed Leiothrix
<i>Alcippe morrisonia</i>	Grey-cheeked Fulvetta
<i>Yuhina castaniceps</i>	Striated Yuhina
<i>Yuhina zantholeuca</i>	White-bellied Yuhina
<i>Parus major</i>	Great Tit
<i>Nectarinia jugularis</i>	Olive-backed Sunbird
<i>Aethopyga christinae</i>	Fork-tailed Sunbird
<i>Dicaeum ignipectus</i>	Fire-breasted Flowerpecker
<i>Dicaeum cruentatum</i>	Scarlet-backed Flowerpecker
<i>Dicaeum concolor</i>	Plain Flowerpecker
<i>Zosterops japonicus</i>	Japanese White-eye
<i>Lanius schach</i>	Long-tailed Shrike
<i>Dicrurus macrocercus</i>	Black Drongo
<i>Dicrurus aeneus</i>	Bronzed Drongo
<i>Urocissa whiteheadi</i>	White-winged Magpie
<i>Dendrocitta formosae</i>	Grey Treepie
<i>Passer montanus</i>	Eurasian Tree Sparrow
<i>Lonchura striata</i>	White-rumped Munia
<i>Lonchura punctulata</i>	Scaly-breasted Munia
<i>Melophus lathami</i>	Crested Bunting

- Oriental Honey Buzzard *Pernis ptilorhynchus*, Crested Serpent Eagle *Spilornis cheela*, Black Kite *Milvus migrans*, Crested Goshawk *Accipiter trivirgatus*, Shikra *Accipiter badius*, Japanese Sparrowhawk *Accipiter gularis*, Besra *Accipiter virgatus*, Eurasian Hobby *Falco subbuteo*, Silver

Pheasant *Lophura nycthemera*, Mountain Scops Owl *Otus spilocephalus*, Collared Scops Owl *Otus bakkamoena*, Oriental Scops Owl *Otus sunia* and Asian Barred Owlet *Glaucidium cuculoides* are Class II Protected nationally.

Reptiles and Amphibians

- Twenty-two species of amphibian and fifteen species of reptile (one turtle, six lizards and eight snakes) were recorded at Shiwandashan during these and earlier surveys (Table 5).
- The identity of *Rana spinulosa*, *Tropidophorus hainanus* and *Amphiesma popei* could not be confirmed. Tadpoles belonging to the genera *Megophrys* and *Leptobrachium* were found but they could not be identified to the exact species due to the lack of adult materials.
- Several specimens of *Geoemyda spengleri*, a globally Endangered and National Class II Protected species, were for sale in the stalls at the Forest Park. They were reported to have been collected (illegally) inside the Park.
- Two additional snake species were observed: *Oligodon formosanus* and *Trimeresurus albolabris*.

Table 5. Amphibians and reptiles recorded in Shiwandashan National Nature reserve and surrounding areas in May 1997, March, April and September 2000. Sequence follows Zhao E.-M. & Adler (1993).

Species	Habitat	May 1997	May 1997	Mar, Apr 2000	Sep 2000
AMPHIBIA					
<i>Megophrys</i> sp.	stream				tadpoles
<i>Leptobrachium</i> sp.	stream				tadpoles
<i>Bufo melanostictus</i>	plantation	✓			
	forest		✓		
	abandoned field				✓
<i>Hyla simplex</i>	pool		✓		
	plantation	✓			
<i>Amolops ricketti</i>	stream		✓		✓
<i>Paa spinosa</i>	stream		✓		
<i>Rana guentheri</i>	plantation	✓			
	marsh		✓		✓
	abandoned field				✓
<i>Rana limnocharis</i>	plantation	✓			
	marsh		✓	✓	
	pool				✓
<i>Rana livida</i>	stream		✓	✓	✓
	forest			✓	
<i>Rana rugulosa</i>	field		✓		
<i>Rana spinulosa</i> ?	forest/stream		✓		
	forest				✓
<i>Rana taipehensis</i>	marsh		✓		
<i>Rana versabilis</i>	riparian forest				✓
	stream				✓
<i>Chirixalus vittatus</i>	pool		✓	✓	
<i>Philautus odontotarsus</i>	forest pool		✓, tadpoles		
<i>Polypedates dennysi</i>	pool		✓		
	forest				✓
<i>Polypedates megacephalus</i>	pool	✓	✓		
<i>Polypedates mutus</i>	pool		✓		
<i>Kalophrynus interlineatus</i>	marsh		✓		
<i>Microhyla butleri</i>	Pool		✓		
	marsh				✓
<i>Microhyla ornata</i>	pool		✓		
<i>Microhyla pulchra</i>	field/pool		✓		
	marsh			✓	
REPTILIA					
<i>Sacalia quadriocellata</i>	stream		✓		
<i>Acanthosaura lepidogaster</i>	forest		✓		✓
<i>Calotes versicolor</i>	plantation	✓			✓
<i>Draco maculatus</i>	fir plantation				✓
<i>Sphenomorphus indicus</i>	forest			✓	✓

<i>Tropidophorus hainanus</i> ?	seepage			✓
<i>Tropidophorus sinicus</i>	forest			✓
	stream			✓
<i>Ahaetulla prasina</i>	forest edge	✓		
<i>Amphiesma popei</i> ?	forest			✓
<i>Calamaria septentrionalis</i>	forest	✓		
<i>Opisthotropis lateralis</i>	stream	✓		
<i>Psammodynastes pulverulentus</i>	forest	✓		
	stream			✓
<i>Ptyas korros</i>	ag. field		✓	
<i>Sinonatrix aequifasciata</i>	stream	✓		✓
<i>Bungarus multicinctus</i>	fir plantation			✓

- Species of conservation concern recorded are:
 - The record of *Leptobrachium* is the first for Guangxi.
 - *Chirixalus vaittatus* has a restricted and fragmented distribution in southern China.
 - *Sacalia quadriocellata* is a globally Endangered species.
- The high diversity of forest and forest stream herpetofauna indicates that some of the forests at Shiwandashan were intact.

Fish

- A total of 18 freshwater fish species were recorded from Shiwandashan National Nature Reserve and Shiwandashan National Forest Park in the September 2000 survey (Table 6).
- Generally, fish abundance was high in the streams visited, especially the forest stream in the Forest Park.
- A number of species are rarely recorded during KFBG's surveys: *Rasbora steineri* is globally restricted to northern Indochina, while *Leiocassis argentivittatus* has so far only been found at Shiwandashan during our surveys.
- Three species could not be identified using existing keys for Chinese freshwater fish; they belong to the Culterinae, Balitoridae and Odontobutidae. They may be of conservation/scientific interest.

Table 6. Freshwater fish recorded from Shiwandashan National Nature Reserve and surrounding area, Southwest Guangxi, 25-27 September 2000. Sequence of families follows Nelson (1994).

Species
<i>Rasbora steineri</i>
<i>Parazacco spilurus spilurus</i>
<i>Zacco platypus</i>
<i>Opsariichthys bidens</i>
<i>Yaoshanicus arcus</i>
<i>Nicholsicypris normalis</i>
Cultrinae sp.
<i>Carassius auratus</i>
<i>Cobitis sinensis</i>
<i>Misgurnus anguillicaudatus</i>
<i>Vanmanenia</i> (cf. <i>caldwelli</i>) sp.
<i>Schistura fasciolata</i>
<i>Leiocassis argentivittatus</i>
<i>Silurus asotus</i>
<i>Pterocryptis gilberti</i>
<i>Mastacembelus armatus</i>
Odontobutidae sp.
<i>Micropercops compressocephalus</i>

- A recent detailed fish survey yielded a total of 102 fish species from the Shiwandanshan area (Zhao & Zhang, 2001). Reported species of particular interest are: *Parazacco spilurus fasciatus*, *Acrossocheilus iridescens*, *Plecoglossus altivelis*, *Botia robusta*, *B. pulchra*, *Cobitis arenae*,

Pseudohemiculter hainanensis, *Rasborinus formosae*, *Xenocypris microlepis*, *Cyprinus acutidorsalis*, *Beaufortia leveretti*, *Cranoglanis boudierius* and *Mastacembelus aculeatus*.

- Together the two surveys recorded over 105 species of freshwater fish from Shiwandashan. This is an exceptionally high figure and the area is of very high conservation value for fish fauna of northern Indochina.

Ants

- Ants were not covered by the present surveys in 2000-2001, but results of an earlier survey of the Forest Park area in May 1997 (Fellowes & Hau, 1997) can be updated here following improved understanding of ant taxonomy (Table 7). At least 65 species were recorded, of which many cannot be firmly identified; most species were recorded inside the National Forest Park.
- The most frequently encountered species included *Crematogaster* sp. 2, *Odontoponera* sp. 1 and *Pheidole smythiesi*.

Table 7. Ant species recorded in and around Shiwandashan National Forest Park, May 1997. “*” = Species with a strong forest association.

Scientific name
<i>Aenictus</i> (<i>ceylonicus</i> group) sp. 1
<i>Aenictus binghami</i> *
<i>Aphaenogaster</i> (cf. <i>beccarii</i>) sp. 1 *
<i>Aphaenogaster</i> (cf. <i>hunanensis</i>) sp. 3 *
<i>Camponotus</i> (cf. <i>anningensis</i>) sp. 39 *
<i>Camponotus</i> (cf. <i>jianghuaensis</i>) sp. 15
<i>Camponotus</i> (cf. <i>mitis</i>) sp. 11
<i>Camponotus</i> (cf. <i>wasmanni</i>) sp. 35
<i>Camponotus</i> (nr. <i>vitreus praerufus</i>) sp. 32
<i>Camponotus rufoglaucus</i> (recorded as <i>Camponotus</i> sp. 22)
<i>Cataulacus granulatus</i>
<i>Cerapachys sulcinodis</i> *
<i>Crematogaster</i> (cf. <i>biroi</i>) sp. 4
<i>Crematogaster</i> (cf. <i>dohrni</i>) sp. 8
<i>Crematogaster</i> (cf. <i>travancorensis</i>) sp. 2
<i>Diacamma</i> (nr. <i>rugosum</i>) sp. 1 (previously recorded as <i>D. pallidum</i>)
<i>Dolichoderus</i> (cf. <i>flatidorsus</i>) sp. 6 (misidentified as <i>Dolichoderus</i> sp. 7 and sp. 8)
<i>Dolichoderus</i> (nr. <i>taprobanae</i>) sp. 4 (misidentified as <i>Dolichoderus</i> (nr. <i>thoracicus</i>) sp. 3)
<i>Hypoponera</i> (cf. <i>excoecata</i>) sp. 2 *
<i>Lepisiota rothneyi</i>
<i>Leptogenys binghamii</i> *
<i>Leptogenys kitteli</i> *
<i>Leptogenys peuqueti</i> (recorded as <i>Leptogenys</i> sp. 13)
<i>Leptogenys</i> (cf. <i>kraepelini</i>) sp. 7 *
<i>Myrmecaria</i> (cf. <i>brunnea</i>) sp. 2
<i>Odontomachus</i> (cf. <i>silvestrii</i>) sp. 2 (misidentified as <i>O. rixosus</i>)
<i>Odontomachus monticola</i> *
<i>Odontoponera</i> (cf. <i>denticulata</i>) sp. 1
<i>Pachycondyla</i> (cf. <i>sharpi</i>) sp. 12
<i>Pachycondyla</i> (<i>javana</i> group) sp. 1 *
<i>Pachycondyla leeuwenhoekii</i> *
<i>Pachycondyla</i> (cf. <i>luteipes</i>) sp. 2 *
<i>Pachycondyla rufipes</i>
<i>Paratrechina</i> (cf. <i>bourbonica</i>) sp. 4
<i>Paratrechina</i> (cf. <i>opaca</i>) sp. 26 (includes specimens recorded as <i>Paratrechina</i> sp. 27) *
<i>Paratrechina</i> (nr. <i>indica</i>) sp. 9 *
<i>Paratrechina longicornis</i>
<i>Paratrechina sauteri</i>
<i>Pheidole</i> (cf. <i>noda</i>) sp. 1
<i>Pheidole nodifera</i>
<i>Pheidole</i> (<i>rinae</i> group) sp. 9
<i>Pheidole smythiesi</i>
<i>Pheidole</i> sp. 11
<i>Pheidole</i> sp. 13A (previously recorded as <i>Pheidole</i> sp. 27) *
<i>Pheidole</i> sp. 28

Scientific name

Pheidologeton sp. 8 *
Polyrhachis (cf. *phalerata*) sp. 2 *
Polyrhachis demangei
Polyrhachis dives
Polyrhachis halidayi
Polyrhachis latona
Polyrhachis tyrannica
Polyrhachis wolffi *
Ponera sp. 5
Prenolepis (cf. *emmae*) sp. 1 *
Pristomyrmex pungens
Pseudolasius sp.
Pyramica dohertyi (recorded as *Smithistruma* sp. 3)
Tapinoma (nr. *indicum*) sp. 2 *
Tapinoma sp. 1
Technomyrmex albipes
Tetramorium sp. 1-group sp. (recorded as *Tetramorium* sp. 14)
Tetramorium (cf. *kraepelini*) sp. 4 *
Tetraoponera binghami (misidentified as *T. attenuata*)
Vollenhovia (cf. *emeryi*) sp. 1 *

- *Ponera* sp. 5 is known only from Shiwandashan.
- Excluding the unique species, 36% of species recorded are forest-associated, a figure typical of secondary forest areas. The proportion was higher (47%) in the more natural vegetation inside the Forest Park, and very low (7%) in the plantation forest outside.

Dragonflies

- Only four species were recorded during the September 2000 survey, probably due to the adverse weather and the lateness of the season (Table 8).

Table 8. Dragonfly species recorded at Shiwandashan National Nature Reserve, September 2000. Sequence follows Schorr *et al.* (2001a, 2001b).

Species
<i>Megalestes</i> sp. n.
<i>Planaeschna risi risi</i>
<i>Orthetrum sabina</i>
<i>Trithemis aurora</i>

- *Megalestes* sp. nov. is different to the undescribed *Megalestes* collected in Damingshan in 2000 and Xidamingshan in 1998; to date it has not been found elsewhere. This species and the aeshnid *Planaeschna risi risi* are forest specialists.
- Many more dragonfly species were recorded during the KFBG survey of this reserve conducted in May 1997 (Fellowes & Hau, 1997), including numerous forest-associated species, listed below: *Philoganga robusta robusta*, *Matrona basilaris basilaris*, *Euphaea guerini*, *Euphaea decorata*, *Rhinocypha perforata perforata*, *Schmidtiphaea vietnamensis*, *Gynacantha japonica*, *Coelliccia cyanomelas*, *Cephalaeschna* sp., *Leptogomphus* sp., *Gomphidia krugeri fukienensis*, *Macromidia rapida*, *Macromia clio*, *Macromia fulgidifrons* (type locality, K.D.P. Wilson, 1998), and *Macromia moorei malayana*.

Butterflies

- Only fourteen butterfly species were encountered during the September 2000 survey (Table 9). This reflects the rainy weather condition encountered that makes the butterflies inactive.
- One species in the genus *Lexias* could not be firmly identified due to the failure to collect a specimen.

Table 9. Butterfly species recorded at Shiwandashan, 25-27 September 2000. Sequence of families follows Bascombe (1995).

Species	Habitat
<i>Lamproptera curia</i>	forest edge
<i>Papilio helenus</i>	forest edge
<i>Papilio paris</i>	forest edge
<i>Papilio polytes</i>	forest edge
<i>Papilio protenor</i>	forest
<i>Troides helena</i>	forest
<i>Eurema hecabe</i>	field
<i>Hebomoia glaucippe</i>	forest edge
<i>Ixias pyrene</i>	forest edge
<i>Abisara echerius</i>	forest edge
<i>Athyma selenophora</i>	forest edge
<i>Cyrestis thyodamus</i>	forest
<i>Euploea mulciber</i>	forest edge
<i>Lexias</i> sp.	forest

- In addition, the following species were recorded during a survey in May 1997 (Fellowes & Hau, 1997): *Abisara neophron*, *Acytolepis puspa*, *Aemona amathusia*, *Amblypodia anita*, *Appias albina*, *Appias lalage*, *Argyreus hyperbius*, *Artogeia canidia*, *Artogeia rapae*, *Astictopterus jama*, *Athyma opalina*, *Athyma perius*, *Atrophaneura varuna*, *Catopsilia pomona*, *Charaxes bernardus*, *Cirrochroa tyche*, *Danaus genutia*, *Delais pasithoe*, *Euploea core*, *Euploea midamus*, *Eurema blanda*, *Eurema laeta*, *Euthalia niepelti*, *Gerosis phisara*, *Graphium agamemnon*, *Halpe* sp., *Heliophorus* sp., *Ideopsis similis*, *Jamides bochus*, *Junonia almana*, *Kankiska canace*, *Lethe confusa*, *Lethe lanaris*, *Lethe naga*, *Lethe sinorix*, *Lethe syrcis*, *Lexias pardalis*, *Limenitis sulphitia*, *Melanitis leda*, *Mooreana trichoneura*, *Neptis hylas*, *Pantoporia hordonia*, *Parantica melanues*, *Parnara bada*, *Pathysa antiphates*, *Polyura eudamippus*, *Prioneris thestylis*, *Spindasis syama*, *Udara albocaerulea* and *Zemeros flegyas*.
- Several forest species were present: *Aemona amathusia*, *Astictopterus jama*, *Euthalia niepelti*, *Lexias pardalis*, *Limenitis sulphitia* and *Mooreana trichoneura*, indicating some good forests are present in Shiwandashan.

Summary of flora and fauna

- The Shiwandashan area has extensive but fragmented secondary northern tropical monsoon forest in a matrix of degraded vegetation and agricultural land. The present surveys recorded 517 vascular plant species, including nine globally Threatened or nationally Protected species. The presence of a number of tropical genera and species and 14 globally restricted species also suggest that the flora is rather distinct from forest vegetation elsewhere in Guangxi. One orchid found, *Bulbophyllum* sp., is new to China. The large number of new distribution records suggests further surveys may reveal findings of conservation importance.
- While Shiwandashan has apparently lost the largest carnivores (such as Tiger *Panthera tigris*, Leopard *P. pardus* and Grey Wolf *Canis lupus*) this extensive mountain range apparently still supports mammals of conservation importance, including the globally Vulnerable Clouded Leopard, Asiatic Golden Cat and Chinese Goral. Some arboreal mammals dependent on mature forest (e.g. Black Giant Squirrel and flying squirrels) also appear to be present.
- One hundred and eleven bird species have been recorded in the surveys at Shiwandashan, including 13 nationally Protected species (mostly raptors and owls). This rather high total was in part due to the influx of migrants in spring and autumn.
- Twenty-two species of amphibian and fifteen species of reptile were recorded at Shiwandashan, including rare species such as the frogs *Leptobranchium* sp. and *Chirixalus vittatus*, and the globally

Endangered turtle *Sacalia quadriocellata* (and probably *Geoemyda spengleri*). A high proportion of species were forest and forest stream specialists.

- From the survey the fish fauna appeared to be of high diversity and abundance, and this impression was confirmed by the notable results of an in-depth survey by Zhao and Zhang (2001).
- Due to the bad weather, the insect fauna was under-recorded during the present surveys. Nevertheless one undescribed damselfly was found. Previous survey in 1997 showed that Shiwandashan had a rich insect fauna that included a number of rare odonates (Fellowes & Hau, 1997).
- Shiwandashan was considered by MacKinnon *et al.* (1996) to be of national biodiversity significance. The present findings confirm the importance of Shiwandashan, which has also been recognised in the Central Government's approval of national-level nature reserve status.

Threats and problems

- Illegal electrofishing was observed in the streams at two locations, affecting the abundance of fish in these sites. In the Forest Park, fishing in the stream was banned, and at the time of the visits fish abundance was much higher there. However, Zhao *et al.* (2002) reported the severity of electrofishing and its adverse impacts on the fish fauna of Shiwandashan. For example, they noticed a sharp decline of fish diversity and abundance in the Forest Park stream during a re-survey in November 2001. The situation for freshwater biota at Shiwandashan is therefore disturbing.
- Villagers at Nadang also collected the frogs *Paa spinosa* for sale. This further affects the stream community and reduces food available for predatory species such as White-eared Night Heron.
- Hawkers at the Forest Park sold the Protected turtle *Geoemyda spengleri* and various forest products (such as bamboo shoots and large legume seeds) that were supposedly collected inside the Park. This small trade poses a threat to the Endangered *G. spengleri* and sets a very bad example for ecotourism practices.
- At Taiping, villagers living next to a good forest block relied on *Illicium* plantation for income. Although there had been plans to relocate villagers and plantation away from the forested area a few years ago, the villagers moved back to plant *Illicium* as they could not generate enough income elsewhere.

Opportunities

- Shiwandashan has recently been designated a National Nature Reserve. It has a large reserve size and still has rather extensive but fragmented cover of secondary forest, which could support viable populations of most or all of the species that remain if protected from further disturbances.
- To increase the conservation value of this reserve, clearing or logging of any scale and for any purposes should be prohibited in the core area. Any existing plantation and agriculture inside the core area should be abandoned to allow natural regeneration of vegetation. Expansion of agricultural land and plantation should be prohibited in the buffer zone and experimental zone, and livestock grazing and small-scale sustainable logging should be carefully controlled.
- Reforestation using an assembly of plant species native to the region may be necessary for the heavily degraded or grazed areas and for rehabilitating plantations of *Illicium*.
- Continued survival of globally Threatened species, such as Clouded Leopard and White-eared Night Heron, might be ensured but only if enforcement of the hunting and logging ban is treated as a top priority.
- To relieve the very high pressure on land and wildlife, the management should consider recruiting local villagers as reserve wardens and forestry police.
- Given the importance of the fish fauna (Zhao and Zhang, 2001), and the occurrence of globally Endangered White-eared Night Heron in Shangsi County (Fellowes *et al.*, 2001), any riparian forest disturbance and illegal fishing should be avoided. The reserve management might consider banning

fishing in certain zones or periods, to allow for the recovery of fish stock, and provision of undisturbed riparian forest for the rare birds to reproduce and forage.

- One of the best-preserved forests visited in the present surveys was at Shiwandashan National Forest Park. Although outside the Nature Reserve, this is protected with an aim to attract ecotourists. The Forest Park has good facilities (such as boarding, restaurants and nature trails) and attracts a good number of visitors. The potential of developing this forest park into an outdoor environmental education centre should be fully explored by putting more emphasis into education display and programme. Activities in conflict with conservation, such as selling plants and animals collected inside the park or including wildlife on restaurant menus, should be discontinued. Guidelines for various aspects of ecotourism development are available, e.g. Ceballos-Lascuráin (1996) and China National Committee of the Man-and-the-Biosphere (1998).
- It has been suggested to extend Shiwandashan Nature Reserve to cover other forest in the north, and to reforest denuded hills (MacKinnon et al., 1996). These measures would provide further support for biodiversity. Transboundary cooperation to protect forest along the border with Vietnam would ensure a larger, more effective protected-area system in the northern Indochina ecoregion.

Acknowledgements

The editors wish to thank the Guangxi Forestry Department for their cooperation and assistance, and all participants of the survey teams, including field staff at Shiwandashan National Nature Reserve and National Forest Park. This work was funded by KFBG.

References

- Anon., 1959-2001. *Flora Reipublicae Popularis Sinicae*. Tomus 2-80. Science Press, Beijing. (In Chinese.)
- Anon., 1991. *Flora of Guangxi*. Volume 1. Guangxi Science Technology Press, Nanning. (In Chinese.)
- Anon., 1996-2001. *Flora of China* Vol. 4, 15, 16, 17, 18, and 24. Science Press, Beijing, and Missouri Botanic Garden Press, St. Louis.
- Anon., 2002a. *Flora of China Checklist*. Published on the Internet: <http://mobot.mobot.org/W3T/Search/foc.html> [Accessed 1 May, 2002.]
- Anon., 2002b. *Flora of China Manuscript*. Published on the Internet: <http://flora.huh.harvard.edu/china/> [Accessed 1 May, 2002.]
- Bascombe, M.J. 1995. Check list of the butterflies of South China. *Memoirs of the Hong Kong Natural History Society* 20: 1-206.
- Ceballos-Lascuráin H., 1996. *Tourism, ecotourism, and protected areas: the state of nature-based tourism around the world and guidelines for its development*. IUCN – The World Conservation Union, Gland, Switzerland, 301 pp.
- Chen, S.-C. (ed.), 1999. Angiospermae Monocotyledoneae Orchidaceae (2). *Flora Reipublicae Popularis Sinicae*. Tomus 18. Science Press, Beijing, 463 pp. (In Chinese.)
- China National Committee of the Man-and-the-Biosphere, 1998. *Nature Reserve and Ecotourism*. China Science and Technology Press, Beijing, 110 pp. (In Chinese.)
- Clements, J.F., 2000. *Birds of the World: A Checklist, Fifth Edition*. Ibis Publishing Company, California, 867pp.
- Fellowes, J.R., and Hau, C.-H., 1997. *A faunal survey of nine forest reserves in Tropical South China, with a review of conservation priorities in the region*. Kadoorie Farm and Botanic Garden, Hong Kong, 151 pp.
- Fellowes, J.R., Zhou, F., Lee, K.S., Hau, B.C.H., Lau, M.W.N., Lam, V.W.Y., Young, L. and Hafner, H., 2001. Status update on White-eared Night Heron *Gorsachius magnificus* in South China. *Bird Conservation International* 11, 101-111.

- Forestry Department of Guangxi Zhuang Autonomous Region, 1993. *Guangxi Nature Reserves*. China Forestry Publishing House, Beijing, 187pp. (In Chinese.)
- Guangdong Forestry Department and South China Institute of Endangered Animals, 1987. *Colour Guide of Wildlife in Guangdong Province*. Guangdong Science and Technology Press, Guangzhou, 139 pp. + 300 colour plates. (In Chinese.)
- Guangxi Forestry Survey and Planning Institute (ed.), 2002. *Guangxi Shiwandashan Natural Reserve Master Plan (2002-2010)*. Guangxi Forestry Survey and Planning Institute and Guangxi Shiwandashan Nature Reserve Management Station, Nanning, 106 pp. (In Chinese.)
- Guangxi Forestry Survey and Planning Institute and Shiwandashan Nature Reserve Management Station (eds.), 2002. *The Comprehensive Investigation Report of Guangxi Shiwandashan Natural Reserve*. Guangxi Forestry Survey and Planning Institute and Shiwandashan Nature Reserve Management Station, Nanning, 363 pp. (In Chinese.)
- Hua, W.L. and Yan, Q.W., 1993. *Protected Animals in China*. Shanghai Scientific and Technological Education Publishing House, Shanghai, 618 pp. (In Chinese with English abstract.)
- Inskipp, T., Lindsey, N. and Duckworth, W., 1996. *An Annotated Checklist of the Birds of the Oriental Region*. Oriental Bird Club, Sandy, Bedfordshire, U.K, 294 pp.
- IUCN, 2002. *2002 IUCN Redlist of Threatened Species*. Published on the Internet: <http://www.redlist.org/> [Accessed 1 November, 2002.]
- Kadoorie Farm and Botanic Garden, 2003. *Report of Rapid Biodiversity Assessments at Damingshan National Nature Reserve, Central Guangxi, China, April and September 2000*. South China Forest Biodiversity Survey Report Series No. 34. KFBG, Hong Kong SAR, ii + 27 pp.
- Lang, K.-Y. (ed.), 1999. Angiospermae Monocotyledoneae Orchidaceae (1). *Flora Reipublicae Popularis Sinicae*. Tomus 17. Science Press, Beijing, 551 pp. (In Chinese.)
- MacKinnon, J., Meng, S., Cheung, C., Carey, G., Zhu, X. and Melville, D., 1996. *A Biodiversity Review of China*. World Wide Fund for Nature (WWF) International, WWF China Programme, Hong Kong, 529 pp.
- Nelson, J.S., 1994. *Fishes of the World, 3rd edition*. John Wiley & Sons, New York, 600 pp.
- Schorr, M., Lindeboom, M. and Paulson, D., 2001a. *List of Odonata of the World (Part 1, Zygoptera and Anisozygoptera)*. July 2001 version. Published on the Internet: <http://www.ups.edu/biology/museum/worldodonates.html>
- Schorr, M., Lindeboom, M. and Paulson, D., 2001b. *List of Odonata of the World (Part 2, Anisoptera)*. April 2001 version. Published on the Internet: <http://www.ups.edu/biology/museum/worldanisops.html>
- State Forestry Administration and Ministry of Agriculture, 1999. *State Protection List of Wild Plants*. (In Chinese.)
- The Plant Names Project, 2002. *International Plant Names Index*. Published on the Internet: <http://www.ipni.org/> [Accessed 1 May, 2002.].
- Tsi, Z.-H. (ed.), 1999. Angiospermae Monocotyledoneae Orchidaceae (3). *Flora Reipublicae Popularis Sinicae*. Tomus 19. Science Press, Beijing, 485 pp. (In Chinese.)
- Vogel, E.F. De and Turner, H., 1992. *Orchid Monographs*. Vol. 6, 49-62.
- Wang, X.P. et al. (eds.), in press. *Plants of the Chinese Region – Status Survey and Conservation Action Plan. Appendix 1 Red Lists*. IUCN/SSC China Plant Specialist Group.
- Wilson, D.E. and Cole, F.R., 2000. *Common Names of Mammals of the World*. Smithsonian Institution Press, Washington and London, xiv + 204 pp.
- Wilson, K.D.P., 1998. *Macromias* from Guangxi Province, China with a description of a new species (Anisoptera:

- Corduliidae). *Odonatologica* 27(4): 469-474.
- Wu, H.L., Shao, K.T. and Lai, C.F. (eds.), 1999. *Latin-Chinese Dictionary of Fishes' Names*. Sueichan Press, Taiwan, 1,028 pp. (In Chinese and English.)
- Zhang, W. (ed.), 1998. *China's Biodiversity: A Country Study*. China Environmental Science Press, Beijing, 476 pp.
- Zhang, Y. *et al.* 1997. *Distribution of Mammalian Species in China*. China Forestry Publishing House, Beijing, 280 pp. (In Chinese and English.)
- Zhao, E., Chang, H.W., Zhao, H. and Adler, K., 2000. Revised Checklist of Chinese Amphibia & Reptilia. *Sichuan Journal of Zoology* 19(3): 196-207. (In Chinese.)
- Zhao, E.-M. and K. Adler, 1993. *Herpetology of China*. Society for the Study of Amphibians and Reptiles, Oxford, Ohio, U.S.A., 522 pp.
- Zhao, Y. and Zhang, C., 2001. Fish fauna and zoogeographical analysis of Shi Wan Da Shan Mountains, Guangxi, China. *Biodiversity Science* : 9(4): 336-344. (In Chinese with English abstract.)
- Zhao, Y. Tang, W. and Zhang, C., 2002. Status and Protection of Fish Resources in the Shiwan Dashan Mountains, Guangxi, China. *Chinese Journal of Zoology* 6: 43-47. (In Chinese with English abstract.)
- Zhou, F., 1996. White-eared Night Heron in Guangxi. Conservation fund in action. *Oriental Bird Club Bulletin* 23, 8-9.




Figure 1. Map showing location of Shiwandashan National Nature Reserve and National Forest Park, Southwest Guangxi, China.